

NC Geometry and Discrete Torsion Fractional Branes: I

El Hassan Saidi*

Sektion der Physik Theorie, Muenchen Universitat, Muenchen

TheresienStr 37, D-80333, Muenchen, Deutschland

and

Lab/UFR-High Energy Physics, Department of Physics

Faculty of Science, Av. Ibn Battouta, P.O Box 1014

Rabat, Morocco

February 1, 2008

Abstract

Considering the complex n -dimension Calabi-Yau homogeneous hyper-surfaces $\mathcal{H}_n$ and using algebraic geometry methods, we develop the crossed product algebra method introduced by Berenstein et Leigh in hep-th/0105229 and build the non commutative (NC) geometries for orbifolds $\mathcal{O} = \mathcal{H}_n/\mathbf{Z}_{n+2}^n$ with a discrete torsion matrix $t_{ab} = \exp[\frac{i2\pi}{n+2}(\eta_{ab} - \eta_{ba})]$, $\eta_{ab} \in SL(n, \mathbf{Z})$. We show that the NC manifolds $\mathcal{O}^{(nc)}$ are given by the algebra of functions on the real $(2n+4)$ Fuzzy torus $\mathcal{T}_{\beta_{ij}}^{2(n+2)}$ with deformation parameters $\beta_{ij} = \exp\frac{i2\pi}{n+2}[(\eta_{ab}^{-1} - \eta_{ba}^{-1}) q_i^a q_j^b]$, q_i^a 's being Calabi-Yau charges of $\mathbf{Z}_{n+2}^n$. We develop graph rules to represent $\mathcal{O}^{(nc)}$ by quiver diagrams which become completely reducible at singularities. Generic points in these NC geometries are be represented by polygons with $(n+2)$ vertices linked by $(n+2)$ edges while singular ones are given by $(n+2)$ non connected loops. We study the various singular spaces of quintic orbifolds and analyze the varieties of fractional D branes at singularities as well as the spectrum of massless fields. Explicit solutions for the NC quintic $\mathcal{Q}^{(nc)}$ are derived with details and general results for complex n dimension orbifolds with discrete torsion are presented.

*H-saidi@fsr.ac.ma

Key words: Non Commutative Geometry, Calabi-Yau Orbifolds, Discrete Torsion, Fuzzy Torus, Crossed Product Algebra, Morita equivalence.

Contents

1	Introduction	4
2	General Setting	7
2.1	Z Representation	8
2.1.1	Correspondence Rule: Commutative $\longrightarrow$ Non Commutative	8
2.1.2	Constraint Eqs: Commutative Geometry is in the centre of the NC algebra.	10
2.2	Φ Representation	12
3	Non commutative Quintic $\mathcal{Q}^{nc}$	16
3.1	Commutative Quintic $\mathcal{Q}$	16
3.2	Non Commutative Quintic: Constrained Method	19
4	NC Geometry versus Discrete Torsion	21
4.1	More on Solutions (3.19)	23
4.2	Discrete Torsion Matrix of $\mathbf{Z}_5^3$	25
4.2.1	Free torsion case	26
4.2.2	Discrete Torsion	27
4.2.3	Quiver Diagrams	31
5	CP Algebra	33
5.1	$\mathcal{Q} \boxtimes \mathbf{Z}_5^3$ CP Algebra	33
5.1.1	The Centre $\mathcal{ZA}(\mathcal{Q} \boxtimes \mathcal{A}_G)$	34
5.1.2	Structure of the Fibration	35
5.2	Explicit Expression of the Φ_i 's	36

6	General Solutions	37
6.1	Morphisms between Eqs(5.1) and Eqs(6.1)	38
6.2	Example	42
7	Fractional Branes	43
7.1	Fixed subspaces of $\mathcal{Q}^{[\alpha]}$	43
7.1.1	$\mathbf{CP}^2 \bowtie \mathcal{S}_1$ Fibration	44
7.1.2	$\mathbf{B} \bowtie \mathcal{S}_{12}$ Fibration	45
7.1.3	Other Fibrations	47
7.2	Fractional Branes on $\mathcal{Q}^{[1]}$	48
7.2.1	Example : $\mathcal{Q}^{[1]} \simeq \mathbf{B}_1 \bowtie \mathcal{S}_1$	48
7.2.2	First Results	48
7.3	Fractional Branes on $\mathcal{Q}^{[2]}$	51
7.3.1	Fibration $\mathcal{Q}^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{12}$	51
7.3.2	Fibration $\mathcal{Q}^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{23}$	52
7.3.3	Fibration $\mathcal{Q}^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{31}$	52
8	Discussion and Conclusion	55
9	References	60

1 Introduction

Recently some interest has been given to build non commutative (NC) extensions of Calabi-Yau orbifolds $\mathcal{O}$, with discrete torsion [1-10]. These NC manifolds, which could play some role in the twisted sector of string compactification on orbifolds, go beyond the standard non commutative $\mathbf{R}_\theta^n$ and NC $\mathcal{T}_\theta^k$ torii examples considered in brane physics [11-16]. They offer a manner to resolve non geometric stringy singularities [17] and present a natural framework to study fractional D branes at the singularities as well as their massless field spectrum [18-23]. The basic idea of this construction may be summarized as follows: the usual closed string compact target subspace $\mathcal{O}$, which might be singular that is $\mathcal{O} \sim \mathcal{M}/G$ with $G \subset \mathbf{Z}_m^k$ having a set of fix points, is viewed as the commuting central subalgebra $\mathcal{Z}(\mathcal{A}^{nc})$ of a non commutative algebra $\mathcal{A}^{nc}$ representing $\mathcal{O}^{nc}$, the NC complex manifold extending $\mathcal{O}$. The non commutative algebra $\mathcal{A}^{nc}$ may be thought of as a fibration $\mathbf{B} \bowtie F$ with a base $\mathbf{B} = \mathcal{O}$ and as a fiber $F = \mathcal{A}_G$ given by the algebra of the group of discrete torsion G .

From the algebraic geometry point of view, the NC manifold $\mathcal{O}^{nc}$ is covered by a finite set of holomorphic matrix coordinate patches $\mathcal{U}_{(\alpha)} = \{Z_i^{(\alpha)}; 1 \leq i \leq n \alpha = 1, 2, \dots\}$ and holomorphic transition functions mapping $\mathcal{U}_{(\alpha)}$ to $\mathcal{U}_{(\beta)}$; $\phi_{(\alpha,\beta)}: \mathcal{U}_{(\alpha)} \rightarrow \mathcal{U}_{(\beta)}$. This is equivalent to say that $\mathcal{A}^{nc}$ is covered by a collection of non commutative local algebras $\mathcal{A}_{(\alpha)}^{nc}$, describing the analytic matrix coordinate patches $\mathcal{U}_{(\alpha)}$ of the NC complex manifold, with analytic maps $\phi_{(\alpha,\beta)}$ on how to glue $\mathcal{A}_{(\alpha)}^{nc}$ and $\mathcal{A}_{(\beta)}^{nc}$. The $\mathcal{A}_{(\alpha)}^{nc}$ algebras have centers $\mathcal{Z}_{(\alpha)} = \mathcal{Z}(\mathcal{A}_{(\alpha)}^{nc})$; when glued together give precisely the commutative target space manifold $\mathcal{O}$. In this way a commutative singularity of $\mathcal{O} \simeq \mathcal{Z}(\mathcal{A}^{nc})$ can be made smooth in the non commutative space $\mathcal{A}^{nc}$ [1, 2]. This idea was successfully used to build NC ALE spaces and some realizations of Calabi-Yau threefolds such as the quintic threefolds $\mathcal{Q}$. In this regards it was shown that the non commutative quintic $\mathcal{Q}^{nc}$ extending $\mathcal{Q}$, when expressed in the coordinate patch $Z_5 = I_{id}$, is given by the following special algebra,

$$\begin{aligned}
Z_1 Z_2 &= \alpha Z_2 Z_1, & Z_3 Z_4 &= \beta \gamma Z_4 Z_3, & (a) \\
Z_1 Z_4 &= \beta^{-1} Z_4 Z_1, & Z_2 Z_3 &= \alpha \gamma Z_3 Z_2, & (b) \\
Z_2 Z_4 &= \gamma^{-1} Z_4 Z_2, & Z_1 Z_3 &= \alpha^{-1} \beta Z_3 Z_1, & (c) \\
Z_i Z_5 &= Z_5 Z_i, & i &= 1, 2, 3, 4;
\end{aligned} \tag{1.1}$$

where α, β and γ are fifth roots of the unity, the parameters of the $\mathbf{Z}_5^3$ discrete group. The Z_i 's are the generators of $\mathcal{A}^{nc}$. One of the main features of this non commutative algebra is that its centre $\mathcal{Z}(\mathcal{Q}^{nc})$ coincides exactly with $\mathcal{Q}$, the commutative quintic threefolds. In [2],

a special solution for this algebra using 5×5 matrices has been obtained and in [24] a class of solutions for eqs(1.1) depending on the Calabi-Yau charges of the quintic threefold has been worked out and some partial results regarding higher dimensional Calabi-Yau hypersurfaces were given. But open questions, such as the explicit dependence in the discrete torsion of the orbifold group, the varieties of the fractional D branes at singularities on such kind of Calabi-Yau orbifolds and more generally fractional branes on NC toric manifolds, have not been answered yet.

Moreover and as far eqs (1.1) are concerned, there are some first remarks that one can make regarding the non commutative structure generated from discrete torsion. Let us focus our attention on the first term of the relations (1.1) namely

$$Z_1 Z_2 = \alpha Z_2 Z_1, \quad (1.2)$$

which, for the case of a complex dimension n Calabi-Yau hypersurface with a $\mathbf{Z}_{n+2}$ discrete group, is supplemented equally by the conditions

$$\begin{aligned} \alpha^{n+2} &= 1, \\ Z_1^{n+2} &\sim I_{id}, \\ Z_2^{n+2} &\sim I_{id}. \end{aligned} \quad (1.3)$$

If for a while one forgets about the full set of eqs(1.1) and considers just the above relations, one discovers that they are nothing but the defining eq of a two dimensional fuzzy torus $\mathcal{T}_\alpha^2$ with a NC parameter α . But where this NC fuzzy two torus comes from? What NC Calabi-Yau manifolds have to do with NC fuzzy torii?

In $4d$ $\mathcal{N} = 4$ $U(N)$ super Yang-Mills gauge theory where eqs type $\phi_1 \phi_2 = \alpha \phi_2 \phi_1$ appear as vacuum configurations of strings on orbifolds preserving some amount of supersymmetric charges, one suspects at first sight that the NC $\mathcal{T}_\alpha^2$ might have origin in the toral subalgebra of the $U(N)$ gauge group. But a careful inspection shows however that this is not true; after all NC calabi-Yau manifolds are mathematical objects and in general have nothing to do with super Yang-Mills theory. As we will show throughout this paper, the origin of $\mathcal{T}_\alpha^2$ is due to a nice property of Z_{n+2} orbifolds; one of the two NC cycles of $\mathcal{T}_\alpha^2$ is related to the $\mathbf{Z}_{n+2}$ discrete orbifold group while the other NC cycle is due an extra discrete hidden symmetry of the Calabi-Yau hypersurface orbifold. To avoid confusion with the original group, we will refer in this paper to the hidden invariance to as $\mathbf{Z}_{n+2}^{aut}$. As such the effective orbifold group should be thought as $\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut}$ rather than $\mathbf{Z}_{n+2}$. The $*$ product should not be confused with the star

product of the Moyal plane one currently used in NC QFT a la Seiberg-Witten; the $*$ symbol is introduced here just to underline that the two $\mathbf{Z}_{n+2}$ and $\mathbf{Z}_{n+2}^{aut}$ groups do not commute. In fact, we will show that to each $\mathbf{Z}_{n+2}$ orbifold subsymmetry of the Calabi-Yau hypersurface orbifold $\mathcal{H}_n/\mathbf{Z}_{n+2}^n$ there exist indeed a $\mathbf{Z}_{n+2}^{aut}$ partner so that the full symmetry is $(\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut})^n$. To fix the ideas, if we denote by $\{\omega_i\}$, the $(n+2)$ commuting group elements of $\mathbf{Z}_{n+2}$ that is $\{\omega_i, \omega_i \omega_j = \omega_j \omega_i; \omega_i^{n+2} = I_{id}\} = \{\omega_i = \omega^i; \omega^{n+2} = I_{id}\}$, and $\{P_{\omega^i}\}$ the elements of its representations $\mathcal{D}(\mathbf{Z}_{n+2})$, then the $\mathbf{Z}_{n+2}^{aut}$ symmetry can be thought as a partner of $\mathbf{Z}_{n+2}$ acting by rotating the ω_i elements amongst themselves. In other words $\mathbf{Z}_{n+2}^{aut}$ is just an automorphism group of $\mathbf{Z}_{n+2}$ operating by adjoint actions. One of the remarkable features of $\mathbf{Z}_{n+2}^{aut}$ is that it can be represented on the same space as that of $\mathcal{D}(\mathbf{Z}_{n+2})$ and moreover does not commute with $\mathbf{Z}_{n+2}$; i.e $\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut} \neq \mathbf{Z}_{n+2}^{aut} * \mathbf{Z}_{n+2}$. Furthermore regarding usually the algebra eqs(1.1) and too particularly the typical relation $Z_1 Z_2 = \alpha Z_2 Z_1$ with eqs(1.3), one may ask why not NC geometries beyond the constraint eqs $\alpha^{n+2} = 1$, $Z_1^{n+2} \sim I_{id}$ and $Z_2^{n+2} \sim I_{id}$. After all, the Fuzzy torus appearing in here is just a special realization of the NC torus namely the rational representation; but there exists also an irrational one, involving Powers Riffel projectors [25-27]. We suspect that such NC geometries exist and are associated with manifolds having a continuous NC $U(1) * U_{aut}(1)$ symmetry which we believe to be the kind of invariance that should appear in building NC toric manifolds. In this first paper, we will address the rational case only, that is the situation where the α parameter is such that $\alpha^{n+2} = 1$, by studying non commutative extensions of orbifolds of complex dimension n Calabi-Yau hypersurfaces with $\widehat{G}$ symmetries contained in $(\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut})^{\otimes n}$. To do so we will need to go beyond the constrained method initiated in [24] and use the algebraic geometry method introduced in [17] which we will first develop it for the case of compact Calabi-Yau manifolds; then we show that this method is a general one that contain the constrained one. After that we apply the new method to build the NC extensions for the large class of Calabi-Yau hypersurfaces with explicit computations for the example of the quintic. Next, using fibration ideas, we develop a way to get the singular spaces and explore different kind of orbifolds and study the various fractional D branes at their singularities as well as their massless field spectrum. In this paper, we shall also give explicit realizations of discrete torsion using the automorphism symmetries of the orbifold group and works out different classes of Morita equivalent solutions for the NC geometry with torsion.

The organization of this paper is as follows: In section 2, we give the general setting for non commutative geometry from discrete torsion by using algebraic geometry techniques. We first develop the constrained method used in [24] and give the extension to higher dimensional

Calabi-Yau hypersurface orbifolds. Then we discuss an other realization using results on $4d$ supersymmetric gauge theories where NC geometry appears as the moduli space of vacuum configurations for superpotentials of type $W(\Phi) = g\varepsilon^{[IJK]} \Phi_I \Phi_J \Phi_K$. In section 3, we give some general features of the quintic symmetries and study its non commutative extensions using the constrained method. Here we also complete partial results obtained in [24]. The analysis developed here applies as well to all Calabi-Yau hypersurfaces. In section 4, we develop non commutative geometry versus discrete torsion and work out the full class of torsion dependent solutions. In section 5, we develop the crossed product (CP) algebra method introduced in [17] and use this development to derive general solutions for the NC Calabi-Yau orbifolds. This analysis recovers the results of [17] as special cases. In section 6, we study the symmetries of the moduli space solutions we have obtained and in section 7, we discuss the varieties of fractional D branes and give the full spectrum of massless fields on the D branes. In section 8, we collect some general results on NC complex d -dimensional Calabi-Yau orbifolds with a discrete torsion matrix $t_{ab} = \exp[\frac{i2\pi}{d+2}(\eta_{ab} - \eta_{ba})]$, $\eta_{ab} \in SL(d, \mathbf{Z})$ and study the classes of varieties of fractional D branes at singularities. The NC manifolds are given by the algebra of functions on the Fuzzy torus $\mathcal{T}_{\beta_{ab}}^{2d}$, where $\beta_{ab} = \exp\frac{i2\pi}{d+2}[(\eta_{cd}^{-1} - \eta_{dc}^{-1}) q_a^c q_b^d]$. Finally we give a conclusion.

2 General Setting

Given a commutative Calabi-Yau manifold $\mathcal{M}$, in particular that class of orbifolds $\mathcal{O}$ with discrete torsion group G , $G \subseteq \mathbf{Z}_{n+2}^n$, described by complex dimension n commutative holomorphic $(n+2)$ homogeneous hypersurface $\mathcal{H}_{n+2} = \mathcal{P}_{n+2}(z_1, \dots, z_{n+2}) = \lambda^{-n-2} \mathcal{P}_{n+2}(\lambda z_1, \dots, \lambda z_{n+2})$, embedded in the complex projective space $\mathbf{CP}^{n+1}$, one can build its non commutative geometry extension $\mathcal{O}^{nc}$ by using discrete torsion cocycles. In this section, we want to show that in general there are two ways one may follow to achieve this goal: (1) A constrained approach using purely algebraic computations and (2) the CP algebra method based on techniques of algebraic geometry and representation theory of G . For pedagogical reasons, we shall refer here below to the first method as the Z realization because of its direct connection with holomorphic analysis of complex manifolds and to the second one as the Φ representation due to its direct physical origin; The CP algebra describes vacuum configurations of some supersymmetric field theories on orbifolds, . We will show that these two realizations are in fact two special basis of the algebra of functionals on fuzzy torii $\mathcal{T}_{\beta_{ij}}^{2(n+2)}$ and are linked to each other by a Morita morphism. In what follows, we develop the general setting of the methods with

special focus on the basic features of each one of the two realizations; more developments and explicit solutions of both of them will be given in the next sections.

2.1 Z Representation

If we denote by $\mathcal{R}_{n+2}$, the ring of holomorphic and degree $(n+2)$ homogeneous polynomials on $\mathbf{CP}^{n+1}$, namely $\mathcal{P}(z_1, \dots, z_{n+2}) = z_1^{n+2} + \dots + z_{n+2}^{n+2} + a_0 \prod_{i=1}^{n+2} z_i = 0$, and by $\mathcal{R}_{n+2}^{[\alpha]} \equiv \mathcal{R}_{n+2}/G_{[\alpha]}$, the set of equivalent classes of homogeneous polynomials that are invariant under the discrete group $G_{[\alpha]} \subseteq \mathbf{Z}_{n+2}^n$, then one can build the non commutative extension of $\mathcal{R}_{n+2}^{[\alpha]}$ by using $G_{[\alpha]}$ discrete torsion cocycles. There are essentially two key steps in using this constrained construction: (1) A correspondence rule which associate to each complex variable z , a chiral matrix operator Z valued in some $G_{[\alpha]}$ representation and (2) a set of constraint eqs ensuring that the original commutative manifold sit in the centre of the NC space. In what follows, we summarize the main lines of this construction.

2.1.1 Correspondence Rule: Commutative $\longrightarrow$ Non Commutative

Starting from the polynomial $\mathcal{P}(z_1, \dots, z_{n+2})$, we associate to each commutative complex holomorphic variable z_i , a (matrix) chiral operator Z_i valued in some representation $\mathcal{D}(G)$ of the orbifold group G . Formally, if $\{T_{\{a\}}; \quad a = 1, \dots\}$ denotes the set of the elements (generators) of the $\mathcal{D}(G)$ representation satisfying the G group law,

$$\begin{aligned} T_{\{a\}} T_{\{b\}} T_{\{a\}}^{-1} &= t_{\{a,b\}} T_{\{b\}}, \\ T_{\{a\}}^{n+2} &= I_{\mathcal{D}(G)} \end{aligned} \quad (2.1)$$

with a given non trivial torsion matrix t_{ab} , and acting on the z_i variables with eigencharges $\lambda_{i\{a\}}$,

$$\begin{aligned} T_{\{a\}} z_i T_{\{a\}}^{-1} &= \lambda_{i\{a\}} z_i, \\ T_{\{a\}}^{n+2} z_i T_{\{a\}}^{-(n+2)} &= z_i \end{aligned} \quad (2.2)$$

then the correspondence rule Commutative-Non commutative, reads as;

$$\begin{aligned} z_i &\longrightarrow Z_i = \prod_a Z_{ia_i} \\ Z_{ia_i} &= x_i \otimes T_{\{a_i\}} \equiv x_i T_{\{a_i\}}. \end{aligned} \quad (2.3)$$

Here the x_i parameters are $\mathbf{C}$ -numbers neutral under the $T_{\{a\}}$ actions and the $\prod_a$ product is carried on the $T_{\{a_i\}}$. From this correspondence, one sees that the Z_i chiral operators inherit features of the $\mathcal{D}(G)$ group representations. In particular, we have for the commutation relations of the Z_{ia_i} , the same cocycles as those appearing in the commutations of the $T_{\{a_i\}}$'s as shown here below:

$$\begin{aligned} Z_{ia_i} Z_{jb_j} &= t_{\{a_i, b_j\}} Z_{jb_j} Z_{ia_i}, \\ T_{\{a_i\}} T_{\{b_j\}} &= t_{\{a_i, a_j\}} T_{\{b_j\}} T_{\{a_i\}}, \\ Z_i Z_j &= t_{ij} Z_j Z_i, \end{aligned} \tag{2.4}$$

with $t_{ij} = \prod_{a,b} t_{\{a_i, b_j\}}$. These relations show clearly that non commutative geometry, as built above, is induced by discrete torsion and has much to do with fuzzy realization of non commutative torii since the above relations look like the familiar $UV = \lambda VU$ equation of the two dimensional NC torus [11, 27]. We will turn to this special issue in the development; for the moment let us comment briefly the above relations. Eqs(2.1) are general relations which may be exploited in different ways. Just for the simple example where $G = \mathbf{Z}_{n+2}$ where we have one apparent generator $T_1 = P_\omega \equiv P$, one can build various torsion matrices $\mathbf{t} = (t_{ab})$. Besides the usual $t_{11} = 1$ term, one can build several non trivial ones. Taking into account the $\mathbf{Z}_{n+2}^{aut}$ automorphism group generated by Q and setting $T_2 = Q_\omega = Q$, the $\mathbf{t}$ matrix reads as

$$t_{ab} = \begin{pmatrix} 1 & \omega^{-1} \\ \omega & 1 \end{pmatrix}. \tag{2.5}$$

An other example involving the $\widehat{\mathbf{Z}_{n+2}} \equiv \mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut}$ group, is to take the T_a operators as given by elements of $\mathbf{Z}_{n+2}$ and $\mathbf{Z}_{n+2}^{aut}$. Generic examples correspond to take T_a as given by r elements of $\mathbf{Z}_{n+2}$ and m elements of $\mathbf{Z}_{n+2}^{aut}$; i.e $\{T_a\} = \{P_{\omega_1}, \dots, P_{\omega_r}; Q_{\omega_1}, \dots, Q_{\omega_m}\}$. In this case torsion is carried by a $(r+m) \times (r+m)$ matrix given by,

$$t_{ab} = \begin{pmatrix} (Id)_{r \times r} & (A)_{r \times m} \\ (B)_{m \times r} & (Id)_{m \times m} \end{pmatrix}, \tag{2.6}$$

where the rectangular A and B matrices read as,

$$A = \begin{pmatrix} \omega^{-1} & \omega^{-1} & \dots & \omega^{-1} & \omega^{-1} \\ \omega^{-2} & \omega^{-2} & \dots & \omega^{-2} & \omega^{-2} \\ \vdots & \vdots & \dots & \vdots & \vdots \\ \omega^{1-m} & \omega^{1-m} & \dots & \omega^{1-m} & \omega^{1-m} \\ \omega^{-m} & \omega^{-m} & \dots & \omega^{-m} & \omega^{-m} \end{pmatrix}_{r \times m},$$

(2.7)

$$B = \begin{pmatrix} \omega & \omega^2 & \dots & \omega^{m-1} & \omega^m \\ \omega & \omega^2 & \dots & \omega^{m-1} & \omega^m \\ \cdot & \cdot & \dots & \cdot & \cdot \\ \cdot & \cdot & \dots & \cdot & \cdot \\ \omega & \omega^2 & \dots & \omega^{m-1} & \omega^m \\ \omega & \omega^2 & \dots & \omega^{m-1} & \omega^m \end{pmatrix}_{m \times r}.$$

It is this kind of matrix torsion which has been used in [24] to build a class of solutions for the NC quintic. Later on we will use an other kind of torsion matrix by focusing our attention on the basic P and Q generators of the $\widehat{\mathbf{Z}_{n+2}^k} \equiv (\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut})^{\otimes k}$ group. To avoid confusion, we will refer to the basic k generators P_ω of $\mathbf{Z}_{n+2}^k$ (respectively Q_ω of $(\mathbf{Z}_{n+2}^{aut})^k$) as E_a , $a = 1, \dots, k$ (respectively J_a).

Concerning eqs(2.3), it is interesting to note that in addition to the $\mathbf{Z}_{n+2}^n$ symmetries eqs(2.2), the z_i complex variables carry equally a $U(1)$ charge of the complex space describing rotations in $\mathbf{R}^2 \simeq \mathbf{C}$ and acting as $z \longrightarrow z \exp i\chi$, with χ a continuous parameter. The splitting eq(2.3) consists to separate the $\mathbf{Z}_{n+2}^n$ and the $U(1)$ charges. The x_i variables carry now the $U(1)$ charge only while the $\mathbf{Z}_{n+2}^n$ charges are carried by the T_a 's; that is,

$$\begin{aligned} T_{\{a\}} x_i T_{\{a\}}^{-1} &= x_i, \\ T_{\{a\}} T_b T_{\{a\}}^{-1} &= \lambda_{i\{b\}} T_b. \end{aligned} \quad (2.8)$$

In practice, and as we will use in section 6, we shall often isolate only the $G_{[\alpha]} = \mathbf{Z}_{n+2}^\alpha$ charges of interest; that is the charges of orbifold subgroup $G_{[\alpha]}$, and so the x_i complex variables still carry the charges of $G/G_{[\alpha]}$. In other words $T_{\{a\}} x_i T_{\{a\}}^{-1} = x_i$ for $T_{\{a\}} \in G_{[\alpha]}$ and $T_{\{a\}} x_i T_{\{a\}}^{-1} = \lambda_{i\{a\}} x_i$ for $T_{\{a\}} \in G/G_{[\alpha]}$.

2.1.2 Constraint Eqs: Commutative Geometry is in the centre of the NC algebra.

Eq(2.4) tells us that the Z_{ia_i} chiral matrix operators may be viewed as the NC generators of a functional algebra $\mathcal{A}_{nc}[Z]$,

$$\mathcal{A}_{nc}[Z] \equiv \mathcal{A}[Z_{1a_1}, \dots, Z_{(n+2)a_{n+2}}], \quad (2.9)$$

living on a commutative orbifold parameterized by the x_i complex variables, but valued in the algebra $\mathcal{A}_{\mathcal{D}(G)}$ of the group representation $\mathcal{D}(G)$. In other words $\mathcal{A}_{nc}[Z]$ may be thought of

as a fiber bundle $\mathcal{B} \boxtimes \mathcal{F}$ whose base $\mathcal{B}$ is $\mathcal{R}_{n+2}/G_{[\alpha]}$ and its fiber $\mathcal{F}$ is just $\mathcal{A}_{\mathcal{D}(G)}$.

$$\mathcal{A}_{nc}[Z] \sim (\mathcal{R}_{n+2}/G_{[\alpha]}) \boxtimes \mathcal{A}_{\mathcal{D}(G)}. \quad (2.10)$$

The commutative orbifold coincide therefore with the centre $\mathcal{Z}(\mathcal{A}_{nc}[Z])$ of the NC algebra $\mathcal{A}_{nc}[Z]$.

$$\mathcal{Z}(\mathcal{A}_{nc}[Z]) \equiv \mathcal{R}_{n+2}/G_{[\alpha]} \quad (2.11)$$

From this property, it follows the constraint relations

$$\begin{aligned} [Z_{ia_i}, Z_{ja_j}^{n+2}] &= 0, \\ \left[Z_{ia_i}, \prod_{j=1}^{n+2} Z_{ja_j} \right] &= 0. \end{aligned} \quad (2.12)$$

The first constraint eq reflects just the group property carried by the second relations of eqs(2.1) and (2.2) while the second constraint has much to do with the Calabi-Yau condition and turns out to play a central role in working out the solutions of the Z_{ia_i} generators.

Using the correspondence eqs(2.3) and by help of the Schur lemma of group representation theory, one sees that the set $\{T_{\{a_i\}}\}$ of the group representation generators appearing in eq(2.3) and the cocycles $t_{\{a_i, a_j\}}$ of eq(2.4) are determined by finding those $T_{\{a_i\}}$ elements satisfying,

$$\begin{aligned} T_{\{a_i\}}^{n+2} &= I_{\mathcal{D}(G)}, \\ \prod_{j=1}^{n+2} T_{a_j} &= I_{\mathcal{D}(G)}. \end{aligned} \quad (2.13)$$

From this brief presentation, one sees that one can get the non commutative extension of $\mathcal{R}_{n+2}/G$ directly by solving the constraint eqs (2.12-13) without need to introduce powerful tools using more involved analysis on the fibration eqs(2.10). This is a strong point in favor of the constrained method; especially that in the Z realization, non commutative geometry is manifest as shown on eqs(2.4). There are however weak points for this constrained method; features such as the fractional D branes at singularities, the explicit dependence of the deformation parameters into the discrete torsion matrix $\mathbf{t}_{ab}$ and hidden quantum symmetries are not completely manifest. Some special issues of the Z representation has been first considered in [2], and re-examined in [24]. Here we will develop the general setting of the method, work out general solutions exhibiting larger symmetries and give further explicit details for the case of the non commutative quintic and more generally for the $\mathcal{H}_n$ hypersurfaces.

2.2 Φ Representation

Building non commutative extensions of Calabi-Yau orbifolds $\mathcal{O}$ with discrete torsion can be achieved by using a more involved method; the CP algebra considered recently in [17]. To get the idea behind this approach and how it appears in supersymmetric field theory embedded in string theory, let us start from the following special chiral superpotential

$$W(\Phi) = g\varepsilon^{[IJK]} \Phi_I \Phi_J \Phi_K \quad (2.14)$$

of four dimension Super Yang-Mills theories; with g a coupling constant, $\varepsilon^{[IJK]}$ is the usual completely antisymmetric and $\Phi = \phi + \theta\psi + \theta\bar{\theta}F$, the $4d$ chiral superfields with θ and $\bar{\theta}$ the well known $4d$ superspace variables [29].

This $W(\Phi)$ superpotential has a manifest $\mathbf{Z}_k \otimes \mathbf{Z}_k$ symmetry, with discrete torsion, respectively generated by $\mathbf{e}$ and $\mathbf{f}$ operators and acting as,

$$\begin{aligned} \mathbf{e} \Phi_I \mathbf{e}^{-1} &= \Phi_I \exp\left(\frac{i2\pi q_I^e}{k}\right), \\ \mathbf{f} \Phi_I \mathbf{f}^{-1} &= \Phi_I \exp\left(\frac{i2\pi q_I^f}{k}\right), \\ \mathbf{f} \mathbf{e} \mathbf{f}^{-1} &= t_{fe} \mathbf{e}, \quad \mathbf{e}^k = I_{id}, \quad \mathbf{f}^k = I_{id}, \end{aligned} \quad (2.15)$$

where the q_I^e and q_I^f charges are such that;

$$\sum_{I=1}^3 q_I^e = 0, \quad \sum_{I=1}^3 q_I^f = 0. \quad (2.16)$$

The form eq(2.14) of the superpotential is very special; it is intimately connected with the Lie algebra invariant form $tr[A, [B, C]]$ and plays an important role in different supersymmetric theories. First, it appears in $4d$ $\mathcal{N} = 4$ supersymmetric field theory with a $U(N)$ gauge group, containing, in addition to the fermions and the $4d$ gauge field A_μ , six scalars fields transforming in the representation $\underline{\mathbf{6}} \times (\underline{\mathbf{N}} \times \overline{\mathbf{N}})$ of $SO(6) \times U(N) \simeq SU(4) \times U(N)$ symmetry. In language of superfields, this corresponds to interpret Φ_I as $\Phi_i = \sum_{\alpha=1}^{N^2} \Phi_{i\alpha} T^\alpha$, $i = 1, \dots, 6$, the coupling parameter g as the $4d$ $\mathcal{N} = 4$ Yang-Mills coupling constant and finally $\varepsilon^{[IJK]}$ as $f^{\alpha\beta\gamma}$, the completely antisymmetric $SU(N)$ Lie algebra structure constants. Put differently, the superpotential of the $4d$ $\mathcal{N} = 4$ SYM expressed in terms of the $4d$ $\mathcal{N} = 1$ superfields is $W(\Phi) \sim g_{SYM}^{N=4} Tr[\Phi_i, [\Phi_j, \Phi_k]]$.

Eq(2.14) may be also thought of as the superpotential of the pure $4d$ $\mathcal{N} = 2$ $SU(2)$ gauge model where the three Φ_I chiral superfields are taken in the adjoint of $SU(2)$, namely

$\Phi_I = \sum_{i=1}^{N^2} \Phi_i \sigma^i$, with σ^α the usual Pauli matrices, which together with the gauge multiplet form a $4d \mathcal{N} = 2$ $SU(2)$ super Yang-Mills representation. In this case, g is the $4d \mathcal{N} = 2$ Yang-Mills coupling constant $g_{SYM}^{N=2}$ and $\varepsilon^{[IJK]}$ is the usual completely antisymmetric $SU(2)$ structure constants ε^{ijk} . Recall in passing that both the $4d \mathcal{N} = 4$ and $\mathcal{N} = 2$ field models may be viewed as the low energy effective field theory of $10d$ strings compactifications on some large volume Calabi-Yau threefolds, such as the six torus T^6 or again the Calabi-Yau elliptic fibrations $K_3 \rtimes T^2$ and $\mathbf{CP}^1 \rtimes \mathbf{CP}^1 \rtimes T^2$.

An other example where the superpotential $W(\Phi) = g\varepsilon^{[IJK]} \Phi_I \Phi_J \Phi_K$ appears is in the low energy field theory of the $10d$ string on $AdS_5 \times S^5/\mathbf{Z}_3$ near the horizon, with a system of N type IIB $D3$ branes located at the orbifold points of $AdS_5 \times S^5/\Gamma \sim \mathbf{R}^{1,3} \times \mathbf{R}^6/\Gamma$. The orbifold discrete group Γ is in the $SO_R(6) \sim SU_R(4)$ R -symmetry and according to whether $\Gamma = I_{id}$, or $\Gamma \subset SU(2) \subset SU_R(4)$ or again $\Gamma \subset SU(3) \subset SU_R(4)$, one gets either $4d \mathcal{N} = 4$ or $4d \mathcal{N} = 2$ or again $4d \mathcal{N} = 1$ supersymmetric theories respectively [30-31]. In the case where $\Gamma = \mathbf{Z}_3 \subset SU(3)$ for instance, the three sets of complex transverse bosonic coordinates ϕ_I of the N $D3$ branes on $\mathbf{C}^3/\mathbf{Z}_3$, are just the scalar components of chiral superfields Φ_i transforming in the bi-fundamentals of $SU(N)^3$; i.e. $\Phi_1 \sim (N, \overline{N}, 1)$, $\Phi_2 \sim (1, N, \overline{N})$ and $\Phi_3 \sim (\overline{N}, 1, N)$. For details see [32].

The next step in this field analysis is to look for supersymmetric vacuum configurations of the supersymmetric scalar potential $V(\phi)$ induced by eliminating the supersymmetric F auxiliary fields. Upon expressing these auxiliary fields F in terms of the ϕ , the vanishing condition of the scalar potential is $V(\phi) = \sum_{I=1}^3 \left| \frac{\partial W}{\partial \phi_I} \right|^2 = 0$ reads in general as;

$$\sum_{I,J=1}^3 [\phi_J \phi_K - \phi_K \phi_J]^2 = 0. \quad (2.17)$$

For complex three dimension orbifolds $\mathcal{M}/\mathbf{Z}_3 \otimes \mathbf{Z}_3$ with discrete torsion, the above condition, which can also be expressed as $\phi_J \phi_K - \phi_K \phi_J = 0$, have to be supplemented with the symmetries eqs(2.15). Putting all these things together, the supersymmetric vacuum configurations, for string compactifications on $\mathbf{Z}_3 \otimes \mathbf{Z}_3$ orbifolds with discrete torsions, reads then as:

$$\begin{aligned} \phi_J \phi_K &= \phi_K \phi_J \\ \mathbf{e} \phi_j \mathbf{e}^{-1} &= \phi_j \exp\left(\frac{i2\pi q_j^e}{3}\right), \\ \mathbf{f} \phi_j \mathbf{f}^{-1} &= \phi_j \exp\left(\frac{i2\pi q_j^f}{3}\right) \\ \mathbf{f} \mathbf{e} \mathbf{f}^{-1} &= t_{fe} \mathbf{e}. \end{aligned} \quad (2.18)$$

Setting $e_a = (e, f)$ and $\lambda_{ai} = (\exp(\frac{i2\pi q_i^e}{3}); \exp(\frac{i2\pi q_i^f}{3}))$, we can rewrite all above informations on supersymmetric vacua on $\mathbf{Z}_3 \otimes \mathbf{Z}_3$ orbifolds into the following condensed form

$$\begin{aligned}\Phi_i \cdot \Phi_j &= \Phi_i \cdot \Phi_j, \\ e_a \Phi_i &= \lambda_{ai} \Phi_i e_a, \\ e_a e_b &= t_{ab} e_b e_a,\end{aligned}\tag{2.19}$$

where we have used capital Greek letters to represent the bosonic scalar components of the chiral fields; they should not be confused with superfields. Actually these relations generate a non commutative crossed product algebra of $\mathbf{C}^3/\mathbf{Z}_3 \otimes \mathbf{Z}_3$. This CP algebra will be denoted as $\mathcal{O} \boxtimes \mathcal{A}_G$, with $\mathcal{O} = \mathbf{C}^3/G$ and $G = \mathbf{Z}_3^2$; it can be thought of as a fibration of the algebra $\mathcal{A}_G$ of the group G on $\mathcal{O}$. We will refer hereafter to eqs(2.19) as the Φ representation of the non commutative orbifold, $\mathcal{O}^{(nc)} \equiv \mathcal{O} \boxtimes \mathcal{A}_G$, and denote it as $\mathcal{A}[\Phi_1, \Phi_2, \Phi_3; \mathbf{e}_1, \mathbf{e}_2]$ or $\mathcal{A}[\Phi; \mathbf{e}]$ for short. The latter, which is just the algebra of functionals on $\mathcal{O}$, is in fact a remarkable basis of $\mathcal{O} \boxtimes \mathcal{A}_G$; An other different, but equivalent, basis of $\mathcal{O}^{(nc)}$ is that given by the following Z realization to which we will also refer to as $\mathcal{A}[Z_1, Z_2, Z_3; e, f]$ or $\mathcal{A}[Z; \mathbf{e}]$ for short,

$$\begin{aligned}Z_i \cdot Z_j &= \beta_{ij} Z_i \cdot Z_j, \\ \mathbf{e}_a \cdot Z_i &= Z_i \cdot \mathbf{e}_a, \\ \mathbf{e}_a \mathbf{e}_b &= t_{ab} \mathbf{e}_b \mathbf{e}_a.\end{aligned}\tag{2.20}$$

The β_{ij} 's are some parameters such as $\beta_{ij}^3 = 1$; their explicit relation with torsion will be given later on. In these equations, the chiral operators Z_i are related to the Φ_i ones by a $\mathbf{e}_a$ dependent (Morita) morphism ϑ_i ; relating the two descriptions; i.e $Z_i = \vartheta_i(e_a) \Phi_i$. More details on this change of frames will be discussed later on; see section 7.

Extension

Given a complex n -dimension orbifold $\mathcal{O}_n = \mathcal{M}_n/G_m$ with G_m , a dimension m discrete group, the Φ basis, $\mathcal{A}[\Phi_1, \dots, \Phi_n; F_1, \dots, F_m] \equiv \mathcal{A}[\Phi; G_m]$, of the non commutative CP algebra $\mathcal{O}_n \boxtimes \mathcal{A}_{G_m}$ is given by the following generalization of eqs(2.19), namely

$$\begin{aligned}\Phi_i \cdot \Phi_j &= \Phi_i \cdot \Phi_j, \quad i, j = 1, \dots, n, \\ F_a \Phi_i &= \lambda_{ai} \Phi_i F_a, \\ F_a F_b &= t_{ab} F_b F_a, \quad a, b = 1, \dots, m.\end{aligned}\tag{2.21}$$

By an appropriate change of basis $Z_i = \vartheta_i(e_a) \Phi_i$, one can also define the Z representation $\mathcal{A}[Z; G_m]$ of the non commutative CP space $\mathcal{O}_n \boxtimes A_{G_m}$ as the algebra generated by the following relations;

$$\begin{aligned} Z_i Z_j &= \beta_{ij} Z_i Z_j, \quad i, j = 1, \dots, n, \\ F_a Z_i &= Z_i F_a, \\ F_a F_b &= t_{ab} F_b F_a, \quad a, b = 1, \dots, m, \end{aligned} \tag{2.22}$$

where now β_{ij} are non trivial cocycles. The difference between eqs(2.21-22) is just an apparent one; both of the Φ and Z frames are in fact Morita equivalent. The key point in the non similarities between the two representations is that in the Φ frame the non commutative orbifold $\mathcal{O}_n^{(nc)}$ is given by a non trivial fibration of the algebra $\mathcal{A}_G$ on $\mathcal{Z}(\mathcal{A}[\Phi; G_m])$; while this fibration becomes trivial in the Z realization. Both Φ and Z describe the NC geometry extension of the same orbifold though non commutative geometry is non manifest on the Φ_i 's. We shall show later on that the Φ representation has however some remarkable features whose main ones are:

(i) fractional D branes phenomenon is manifest in the Φ realization; it is connected with complete reducibility of the representation at singularities. In other words, we will show that the non zero Φ_i matrix brane transverse coordinates decompose at the orbifold points as follows

$$\Phi_i = \sum_{k=1}^{\dim \mathcal{D}(G_m)} \phi_{ik} \Pi_k \tag{2.23}$$

where Π_k are the projectors on the eigen states basis of the $\mathcal{D}(G_m)$ representation and where ϕ_{ik} are the transverse coordinates of the fractional D branes.

(ii) Along with the commutative G_m symmetry, the NC algebra has an extra automorphism G_m^{aut} embodied by the dual group $\tilde{G}_m$ operating on G_m through the following adjoint action;

$$J_a E_b J_a^{-1} = \alpha_{ab} E_b, \tag{2.24}$$

In this relation, $\{E_a\}$ and $\{J_a\}$ are respectively the generators of the $\mathcal{D}(G_m)$ and $\mathcal{D}(G_m^{aut})$ representations and the complex number α_{ab} are some given group cocycles which are basically given by the G_m group elements. As such the group symmetry of the commutative geometry will be not only given by the G_m group, but rather a larger non commutative one namely $G_m * G_m^{aut}$.

(iii) Solutions for the Φ_i 's can be easily worked out; one has just to take the Φ_i 's either in $\mathcal{D}(G_m)$ or in $\mathcal{D}(G_m^{aut})$. In other words, the Φ_i 's are solved either by help of the E_a generators of $\mathcal{D}(G_m)$ as shown here below; see eqs(4.9-10) for the quintic case,

$$\Phi_i = \phi_i \prod_{a=1}^m E_a^{p_i^a}, \quad (2.25)$$

or again by help of the J_a generators of the automorphism symmetry $\mathcal{D}(G_m^{aut})$ as; see also eqs(5.16),

$$\Phi_i = \phi_i \prod_{a=1}^m J_a^{q_i^a}. \quad (2.26)$$

In this paper we will develop further the CP algebra method for a large class of Calabi-Yau orbifolds with discrete torsion and too particularly orbifolds of the Calabi-Yau homogeneous hypersurfaces $\mathcal{H}_n$. As a toy model, we will focus our attention on the non commutative quintic $\mathcal{Q}$. There, the discrete symmetry group G will be taken as Z_5^3 for $\mathcal{Q}$ and more generally as Z_n^{n+2} for complex n dimensional Calabi-Yau hypersurfaces. NC *non homogeneous* Calabi-Yau hypersurfaces, and too particularly the so called local Calabi-manifolds using toric methods, will be described elsewhere[33]

3 Non commutative Quintic $\mathcal{Q}^{nc}$

For self containment, we deserve this section to recall some general features on the commutative quintic; then we expose the constrained method for deriving the Z realization of the non commutative one; $\mathcal{Q}^{nc}$.

3.1 Commutative Quintic $\mathcal{Q}$

To begin consider the complex analytic homogeneous hypersurface,

$$\mathcal{P}_5(z_1, \dots, z_5) = z_1^5 + z_2^5 + z_3^5 + z_4^5 + z_5^5 + a_0 \prod_{i=1}^5 z_i = 0, \quad (3.1)$$

where $(z_1, z_2, z_3, z_4, z_5)$ are commuting homogeneous coordinates of $\mathbf{CP}^4$, the complex four dimensional projective space, and a_0 a non zero complex moduli parameter. This polynomial describes a well known Calabi-Yau threefolds namely the commutative quintic denoted in

this paper by $\mathcal{Q}$. Besides invariance under permutations of the z_i variables, $\mathcal{Q}$ has a set of geometric discrete isometries acting on the homogeneous coordinates z_i as:

$$z_i \rightarrow z_i \omega^{q_i^a}, \quad (3.2)$$

with $\omega^5 = 1$ and q_i^a integer charges to which we will refer here below as the Calabi-Yau charges. These are the entries of the following five dimensional vectors,

$$\begin{aligned} \mathbf{q}^1 &= (q_1^1, q_2^1, q_3^1, -q_1^1 - q_2^1 - q_3^1, 0) \\ \mathbf{q}^2 &= (q_1^2, q_2^2, q_3^2, -q_1^2 - q_2^2 - q_3^2, 0) \\ \mathbf{q}^3 &= (q_1^3, q_2^3, q_3^3, -q_1^3 - q_2^3 - q_3^3, 0). \end{aligned} \quad (3.3)$$

In these relations, we have set $q_5^a = 0$, a useful feature which can be thought of as corresponding to working in $\mathcal{U}_\alpha[z_1, \dots, z_4, z_5 = 1]$, the $\mathcal{Q}$ local coordinate patch where $z_5 = 1$. For illustrating applications, we will mainly use the following special choice $q_1^a = 1, q_2^1 = q_3^2 = q_4^3 = -1$; all remaining ones are equal to zero. In other words;

$$\begin{aligned} \mathbf{q}^1 &= (1, -1, 0, 0, 0) \\ \mathbf{q}^2 &= (1, 0, -1, 0, 0) \\ \mathbf{q}^3 &= (1, 0, 0, -1, 0). \end{aligned} \quad (3.4)$$

The q_i^a charges in eqs(3.2) and eqs(3.3-4) are defined modulo five; $q_i^a \equiv q_i^a + 5\mathbf{Z}$, and satisfy naturally the identities

$$\sum_{i=1}^5 q_i^a = 0 \quad \text{modulo}(5); \quad a = 1, 2, 3. \quad (3.5)$$

These constraint eqs are necessary as required by invariance under eqs(3.2) of the $a_0 \prod_{i=1}^5 z_i$ monomials of eq(3.1); they ensure that the holomorphic hypersurface(3.1) is a Calabi-Yau manifold. The constraint eqs(3.5) are just the analogue of the vanishing of the first Chern class of the quintic; $c_1(\mathcal{Q}) = 0$, in differential geometry language. These are also the conditions under which the underlying effective field theory flows in the infrared to a CFT [34, 35].

In section 6, we shall show that given the q_i^a charges and a torsion matrix, we can build extra symmetries of eqs(3.1) acting as $z_i \rightarrow z_i \omega^{p_i^a}$ where now the p_i^a charges are given by,

$$\begin{aligned} p_i^a &= \eta_{ab} q_i^b, \\ \sum_{i=1}^5 p_i^a &= 0, \end{aligned} \quad (3.6)$$

with η_{ab} a 3×3 matrix of $SL(3; \mathbf{Z})$. The η_{ab} matrix turns out to play an important role in our present study; it encodes the effects of $G^{aut} \simeq \mathbf{Z}_5^{aut\ 3}$; the group of automorphisms of G . We will show later that the matrix η_{ab} is the carrier of the discrete torsion of the $\mathbf{Z}_5^3$ symmetry; its antisymmetric part $\eta_{ab} - \eta_{ba}$ is related to the logarithm of the t_{ab} matrix of eqs(2.1).

In summary, one should retain that, if we denote by $\mathcal{R}_5[[z_1, \dots, z_5]]$; $\mathcal{R}_5$ for short, the ring of complex holomorphic and homogeneous polynomials of degree five on $\mathbf{CP}^4$ and by $G_{[\nu]} \subseteq G = \mathbf{Z}_5^3$, the subgroups of G and $G_{[\nu]}^{aut} \subseteq G^{aut} \simeq \mathbf{Z}_5^{aut\ 3}$; the automorphisms of $G_{[\nu]}$, then one can build various orbifolds of the commutative quintic as shown here below,

$$\mathcal{Q}^{[\nu]} = \mathcal{R}_5 / G_{[\nu]}. \quad (3.7)$$

For later use, we will consider the two following examples of the quintic orbifolds $\mathcal{Q}^{[1]}$ and $\mathcal{Q}^{[2]}$ associated respectively with $G_{[1]} = \mathbf{Z}_5$ and $G_{[2]} = \mathbf{Z}_5^2$ subgroups of $G = \mathbf{Z}_5^3$,

$$\begin{aligned} \mathcal{Q}^{[1]} &= \mathcal{R}_5 / \mathbf{Z}_5, \\ \mathcal{Q}^{[2]} &= \mathcal{R}_5 / \mathbf{Z}_5^2. \end{aligned} \quad (3.8)$$

Throughout this study, we will mainly stay in the coordinate patch $\mathcal{U}_{(\alpha)}[z_1, \dots, z_4, z_5 = 1]$ as required by the q_i^a choice made in eqs(3.4). The move to an other patch of the quintic, say $\mathcal{U}_{(\beta)}[w_1, \dots, w_4, w_5 = 1]$ with Calabi-Yau charges r_i^a , is ensured by holomorphic transition functions $\phi_{(\alpha, \beta)}^{(r_i^a, q_i^a)}$ carrying appropriate $\mathbf{Z}_5^3$ charges. Note that on the coordinate patch $\mathcal{U}_{(\alpha)}[z_1, \dots, z_4, z_5 = 1]$, the full $\mathbf{Z}_5^3$ orbifold symmetry has no fix point, the only stable one, namely $(0, 0, 0, 0, 1)$, does not belong to $\mathcal{R}_5 / \mathbf{Z}_5^3$. However thinking of the quintic orbifold $\mathcal{R}_5 / \mathbf{Z}_5^3$ as either a $\mathbf{Z}_5$ orbifold of $\mathcal{R}_5 / \mathbf{Z}_5^2$, that is $\mathcal{R}_5 / \mathbf{Z}_5^3 \sim (\mathcal{R}_5 / \mathbf{Z}_5^2) / \mathbf{Z}_5 \sim \mathcal{R}_5' / \mathbf{Z}_5$ or again as a $\mathbf{Z}_5^2$ orbifold of $\mathcal{R}_5 / \mathbf{Z}_5$; i.e $\mathcal{R}_5 / \mathbf{Z}_5^3 \sim (\mathcal{R}_5 / \mathbf{Z}_5) / \mathbf{Z}_5^2 \sim \mathcal{R}_5'' / \mathbf{Z}_5$, one can consider the fixed points of the respective singular spaces of $\mathcal{R}_5' / \mathbf{Z}_5$ and $\mathcal{R}_5'' / \mathbf{Z}_5$. This procedure is also equivalent to set to zero some of the Calabi-Yau vector charges associated with the $\mathbf{Z}_5^3$ symmetry. For example the orbifold $\mathcal{R}_5'' / \mathbf{Z}_5$ may also be thought of as $\mathcal{R}_5 / \mathbf{Z}_5$ by setting $\mathbf{q}^2 = \mathbf{q}^3 = \mathbf{0}$

and so may be compared to $\mathcal{Q}^{[1]}$ while by using a similar reasoning $\mathcal{R}_5'/\mathbf{Z}_5$ can be compared with $\mathcal{Q}^{[2]}$. In section 7, we shall use both of the spaces to study fractional branes and give more explicit examples.

3.2 Non Commutative Quintic: Constrained Method

A way to get non commutative extensions of the quintic $\mathcal{Q}$ by using discrete torsions, is to start from the complex homogeneous hypersurface (3.1) and choose a coordinate patch, say $z_5 = 1$ and so $q_5^a = 0$. Then associate to the $\{z_1, z_2, z_3, z_4\}$ local variables, the set of 5×5 matrix operators $\{Z_1, Z_2, Z_3, Z_4\}$ and Z_5 with the identity matrix I_{id} . The non commutative quintic $\mathcal{Q}^{nc}$, associated to eqs(3.1) and eqs(3.4) is a non commutative algebra generated by the Z_i 's. It is just a special subalgebra of the ring of functions on the space of matrices $\mathcal{M}at(5, \mathbf{C})$ and reads in term of the Z_i generators as;

$$\begin{aligned} Z_i Z_j &= \beta_{ij} Z_j Z_i \quad i, j = 1, \dots, 4, \\ Z_i Z_5 &= Z_5 Z_i, \end{aligned} \tag{3.9}$$

where β_{ij} is an invertible matrix constrained by

$$\begin{aligned} \beta_{ji} &= \beta_{ij}^{-1} \\ \beta_{j1}\beta_{j2}\beta_{j3}\beta_{j4} &= 1, \quad \forall j, \\ \beta_{i1}^5 &= 1, \quad \forall i, \\ \beta_{i2}^5 &= 1, \quad \forall i, \\ \beta_{i3}^5 &= 1, \quad \forall i, \\ \beta_{i4}^5 &= 1, \quad \forall i. \end{aligned} \tag{3.10}$$

These constraint eqs reflect just the property that the commutative $\mathcal{Q}$ should be in the centre $\mathcal{Z}(\mathcal{Q}^{nc})$ of eqs(3.7). In other words $\mathcal{Q} \equiv \mathcal{Z}(\mathcal{Q}^{nc})$, or equivalently;

$$\begin{aligned} [Z_j, Z_i^5] &= 0, \\ \left[Z_j, \prod_{i=1}^4 Z_i \right] &= 0. \end{aligned} \tag{3.11}$$

A class of solutions for possible β_{ij} 's is obtained as follows: First parameterize β_{ij} as;

$$\beta_{ij} = \exp i \left(\frac{2\pi}{5} L_{ij} \right) = \omega^{L_{ij}}. \tag{3.12}$$

where L_{ij} is a 5×5 antisymmetric matrix satisfying moreover

$$\sum_{i=1}^5 L_{ij} = 0. \quad (3.13)$$

A careful inspection of eqs(3.11) shows that this constraint comes from the term $a_0 z_1 z_2 z_3 z_4 z_5$ exactly as for the Calabi-Yau condition eqs(3.5). This means that a possible solution of (3.11) is to take L_{ij} as

$$L_{ij} = n_{ab} (q_i^a q_j^b - q_j^a q_i^b) \equiv m_{ab} q_i^a q_j^b, \quad (3.14)$$

where n_{ab} is a priori an arbitrary 3×3 matrix of integer entries. The matrix L_{ij} is built as bi-linears of the q_k^c charge vectors; ensuring automatically the condition $\sum_{i=1}^5 L_{ij} = 0$. This why, we shall still refer to the constraint $\sum_{i=1}^5 L_{ij} = 0$ as the Calabi-Yau condition. Moreover, since eq(3.14) can also be written as $L_{ij} = m_{ab} q_i^a q_j^b$, with $m_{ab} = n_{ab} - n_{ba}$, one suspects that the m_{ab} matrix should be linked to the structure constants of the underlying $(\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3$ symmetries. We will show, in section 6, that L_{ij} should read in fact as;

$$L_{ij} = p_i^a q_j^a - p_j^a q_i^a, \quad (3.15)$$

where p_i^a are the charges of a non manifest invariance of eqs(3.1). One may think about this latter invariance as a hidden symmetry induced by discrete torsions and, by analogy with $\mathbf{Z}_5^3$, can be defined as acting on the z_i variables as;

$$z_i \rightarrow z_i \omega^{p_i^a}, \quad (3.16)$$

with $p_i^a = n_{ab} q_i^b$. We will prove moreover that the m_{ab} coupling matrix is nothing but the inverse of the logarithm of the discrete torsion matrix $\mathbf{t}_{ab}$ of the group $(\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3$.

To have an idea on how the formulae of the constrained method work in practice, let us go the local patch $z_5 = 1$ with q_i^a charges as in eq(3.4) and perform some explicit calculations. Setting $m_{12} = k_1$, $m_{23} = k_2$ and $m_{13} = k_3$ are integers modulo 5, the L_{ij} matrix reads as:

$$L_{ij} = \begin{pmatrix} 0 & k_1 - k_3 & -k_1 + k_2 & k_3 - k_2 & 0 \\ -k_1 + k_3 & 0 & k_1 & -k_3 & 0 \\ k_1 - k_2 & -k_1 & 0 & k_2 & 0 \\ -k_3 + k_2 & k_3 & -k_2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (3.17)$$

In this case, the generators of the non commutative quintic $\mathcal{Q}^{nc}$ satisfy the following algebra:

$$\begin{aligned}
Z_1 Z_2 &= \omega^{k_1-k_3} Z_2 Z_1, & Z_1 Z_3 &= \omega^{-k_1+k_2} Z_3 Z_1, \\
Z_1 Z_4 &= \omega^{k_3-k_2} Z_4 Z_1, & Z_2 Z_3 &= \omega^{k_1} Z_3 Z_2, \\
Z_2 Z_4 &= \omega^{-k_3} Z_4 Z_2, & Z_3 Z_4 &= \omega^{k_2} Z_4 Z_3.
\end{aligned} \tag{3.18}$$

Putting $\alpha = \omega^{k_1-k_3}$, $\beta = \omega^{k_1-k_2}$ and $\gamma = \omega^{k_3}$, one discovers the algebra of [2] given by eqs(1.1). Moreover the solution of these eqs read, up to a normalization factor, as:

$$\begin{aligned}
Z_1 &= x_1 \mathbf{P}_{\omega^{k_1+k_2+k_3}} \mathbf{Q}^3 \\
Z_2 &= x_2 \mathbf{P}_{\varpi^{k_1}} \mathbf{Q}^{-1} \\
Z_3 &= x_3 \mathbf{P}_{\varpi^{k_2}} \mathbf{Q}^{-1} \\
Z_4 &= x_4 \mathbf{P}_{\varpi^{k_3}} \mathbf{Q}^{-1},
\end{aligned} \tag{3.19}$$

where x_i are as in eqs(2.), ϖ is the complex conjugate of ω and where

$$\mathbf{P}_\alpha = \text{diag}(1, \alpha, \alpha^2, \alpha^3, \alpha^4), \quad \mathbf{Q} = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix} \tag{3.20}$$

with α standing for $\omega^{k_1}, \omega^{k_2}, \omega^{k_3}$ and their products. This solution shows clearly that Z_i^5 , the product $\prod_{i=1}^4 Z_i$ and their linear combination are all of them in the centre $\mathcal{Z}(\mathcal{Q}^{nc})$ of the non commutative algebra $\mathcal{Q}^{nc}$.

4 NC Geometry versus Discrete Torsion

In quantum physics, non commutativity appears in different ways and has various origins and different interpretations [36, 11, 12]. At very low energy effective field theory models such as in the Chern Simons model of the fractional quantum Hall effect [37, 38, 39], NC geometry is generated by a stronger enough constant external magnetic field B which couples the electrons two dimensional position vectors $x^i(t)$ as $B\varepsilon_{ij}x^i(t)\frac{\partial x^j(t)}{\partial t}$. Quantization rules lead to a non vanishing commutator for the position vectors; i.e $[x^i, x^j] = i\frac{\epsilon^{ij}}{B\nu_L}$, with $\nu_L = \frac{1}{k}$ being

the Laughlin filling fraction. For strong B , the quantum properties of the system of electrons are described by a non commutative Chern-Simons gauge theory on a D2 brane. Such a 2D condensed matter phase has received recently an important interest due to similarities with branes systems in superstring theories. At very high energies, say around the Planck scale as in string theory, non commutativity is generated by the antisymmetric NS-NS $B_{\mu\nu}$ field and it is linked to the existence of open strings ending on D branes and a dynamics governed by a boundary conformal invariance [12]. This issue has been subject to much interest during the few last years in connection with the derivation of non commutative solitons and the study of the tachyon condensation by following the GMS method. In M theory, non commutative geometry comes as a non trivial solution for the matrix model compactification, but still generated by an antisymmetric field; the eleven dimensional three form gauge field $C_{\mu\nu\rho}$ [11]. Elements g_1 and g_2 of the group of automorphism symmetries of the matrix model on a two torus $\mathcal{T}^2$ are in general governed by the central relation $g_1 g_2 g_1^{-1} g_2^{-1}$ taken to be proportional to the identity operator; that is $g_1 g_2 g_1^{-1} g_2^{-1} = \lambda I_{id}$, with $\lambda \in \mathbb{C}^*$ as required by the Schur lemma [11, 14]. In quantum mathematics, NC geometry is viewed as an algebraic structure $\mathcal{M}_\hbar[X_1, \dots, X_N]$ going beyond the usual $\mathcal{C}[x_1, \dots, x_N]$ commutative one, with the ideal $\{x_i x_j = x_j x_i\}$. Formally, the generic commutation relations of the generators of the quantum algebra $\mathcal{M}_\hbar[X_1, \dots, X_N]$, may be written as [40, 41]:

$$X_I * X_J = r_{IJ}^{KL}(X) X_K * X_L + b_{IJ}(x), \quad (4.1)$$

where $r_{IJ}^{KL}(X)$ and $b_{IJ}(X)$ are some polynomials in X_I which may be thought of as;

$$\begin{aligned} r_{IJ}^{KL}(X) &= \delta_I^L \delta_J^K + \hbar r'_{IJ}{}^{KL} + \dots, \\ b_{IJ}(X) &= \hbar (\Omega_{IJ} + f_{IJ}^K X_K + \dots) \end{aligned} \quad (4.2)$$

In the limit $\hbar \rightarrow 0$, $r_{IJ}^{KL}(X) \rightarrow \delta_I^L \delta_J^K$, $b_{IJ}(X) \rightarrow 0$; and so one recovers the usual commutative structure of $\mathcal{C}[x_1, \dots, x_N]$. For the general cases, such for instance $(r_{IJ}^{KL}(x), b_{IJ}(x))$ equals to $(0, B_{IJ})$, $(0, f_{IJ}^K X_K)$ and $(R_{IJ}^{KL}, 0)$, one gets respectively the canonical commutator, the Lie algebra bracket and the quantum Yang-Baxter spaces [36].

The non commutative structure we are dealing with in the present paper corresponds to an other special situation where $b_{IJ}(x) = 0$ and,

$$r_{IJ}^{KL}(x) = \beta_{KL} \delta_I^L \delta_J^K, \quad (4.3)$$

with β_{IJ} is a root of unity. This NC geometry is generated by discrete torsion matrix of the orbifold group $\mathbf{Z}_5^3$ and has much to do with NC Fuzzy torii representations. Since discrete

torsions are involved in string compactifications on orbifolds and twisted string sectors, one expects that such NC structure could play some role in string theory on orbifolds. As we shall show by explicit analysis in this paper, see sections 4 and 7, NC geometry versus discrete torsion leads to fractional D branes at orbifold singularities and offers a way to resolve non geometric singularities. Points of the usual geometry are replaced by polygons in the non commutative case.

To get the right link between the non commutative quintic solutions given in the previous section and the discrete torsions of $\mathbf{Z}_5^3$, we start by reconsidering the solutions eqs(3.19) by first showing that they are not so general as claimed in [24]. Then we give some results regarding $\mathbf{Z}_5^3$ group theoretical representations and present a realization for the case where discrete torsions are non trivial. Next, we study the $\mathbf{Z}_5^{aut3}$ hidden symmetry of the quintic raised in the previous sections and acting as in eqs(3.16).

4.1 More on Solutions (3.19)

A careful inspection of eqs(3.19), shows that this solution involves various group elements of the five dimensional representation of $\mathbf{Z}_5 * \mathbf{Z}_5^{aut}$; these are $\mathbf{P}_{\omega^{k_1}}, \mathbf{P}_{\omega^{k_2}}$ and $\mathbf{P}_{\omega^{k_3}}$ and powers of $\mathbf{Q}$. Since $\mathbf{P}_\alpha \mathbf{P}_\beta = \mathbf{P}_{\alpha\beta}$, it follows that these solutions may also be written as

$$\begin{aligned} Z_1 &= x_1 \mathbf{P}^{k_1+k_2+k_3} \mathbf{Q}^3 \\ Z_2 &= x_2 \mathbf{P}^{-k_1} \mathbf{Q}^{-1} \\ Z_3 &= x_3 \mathbf{P}^{-k_2} \mathbf{Q}^{-1} \\ Z_4 &= x_4 \mathbf{P}^{-k_3} \mathbf{Q}^{-1}, \end{aligned} \tag{4.4}$$

where we have set $\mathbf{P}_\omega = \mathbf{P}$. The symmetry of the solutions (3.19) is then $\mathbf{Z}_5 * \mathbf{Z}_5^{aut}$ generated by $\mathbf{P}$ and $\mathbf{Q}$ satisfying $\mathbf{Q} \mathbf{P} \mathbf{Q}^{-1} = \omega \mathbf{P}$. Comparing this relation with eq(2.3), i.e $Z_i = x_i \prod_a T_{\{a_i\}}$, one may identify easily the various elements of $\mathbf{Z}_5 * \mathbf{Z}_5^{aut}$ entering in the building of the solutions. For the operators system $\{\mathbf{P}_{\omega^{k_1}}, \mathbf{P}_{\omega^{k_2}}, \mathbf{P}_{\omega^{k_3}}, \mathbf{Q}\}$ appearing in the building of Z_2, Z_3 and Z_4 , we have the following torsion matrix;

$$t_{\mu\nu} = \begin{pmatrix} 1 & 1 & 1 & \omega^{k_1} \\ 1 & 1 & 1 & \omega^{k_2} \\ 1 & 1 & 1 & \omega^{k_3} \\ \omega^{-k_1} & \omega^{-k_2} & \omega^{-k_3} & 1 \end{pmatrix}. \tag{4.5}$$

Setting $\theta_{\mu\nu} = \frac{-5i}{2\pi} \log \mathbf{t}_{\mu\nu}$, we can rewrite the above relation as

$$\mathbf{t}_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & k_1 \\ 0 & 0 & 0 & k_2 \\ 0 & 0 & 0 & k_3 \\ -k_1 & -k_2 & -k_3 & 0 \end{pmatrix}; \quad (4.6)$$

showing clearly that discrete torsion exist whenever one of the k_a integers are non zero; i.e $k_a \neq 0$. One of the remarks concerning eqs(3.19) is that $(\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3$ invariance is not fully apparent. We will turn to this feature later on when we derive the general solutions later on; but for the moment let us make a remark concerning the NC solutions one gets in general.

Due to discrete torsions, the algebraic structure of the D branes wrapping the compact manifold change. Brane points $\{z_i\}$ of commutative geometry become, in presence of torsion, fibers based on $\{z_i\}$ and valued in the algebra $\mathcal{A}_G$ of the torsion group G . As such, the correspondence;

$$z_i \cdot 1 = \rightarrow Z_i = \sum_{k,l=1}^5 Z_i^{kl} |k \rangle \langle l|. \quad (4.7)$$

has a nice interpretation in terms of quiver diagrams. Associating to each $\mathbf{e}_{kl} \equiv |k \rangle \langle l|$ matrix vector basis, a straight line (string) oriented from the k end to the l one and to each $\mathbf{e}_{kk} = \pi_k \equiv |k \rangle \langle k|$ projector, a loop starting and ending at the position k , as shown on the following table, one may draw a quiver diagram for each Z_i matrix generator of the NC algebra.

Operators	diagrams
$\mathbf{a}_k^+ \equiv k \rangle \langle k+1 $	$k \quad \bullet \longrightarrow \bullet \quad (k+1)$
$\mathbf{a}_k^- \equiv k+1 \rangle \langle k $	$(k+1) \quad \bullet \longleftarrow \bullet \quad k$
$\prod_{j=0}^n \mathbf{a}_{k+j}^+ \equiv k \rangle \langle k+n $	$k \quad \bullet \longrightarrow \bullet \quad (k+n)$
$\prod_{j=0}^n \mathbf{a}_{k+j}^- \equiv k+n \rangle \langle k $	$(k+n) \quad \bullet \longleftarrow \bullet \quad k$
$\pi_k = \mathbf{a}_k^+ \mathbf{a}_k^- = k \rangle \langle k $	$k \bullet \rightleftharpoons \bullet (k+1) \equiv k \bigcirc \equiv \bullet$
$\pi_k = \mathbf{a}_{k-1}^+ \mathbf{a}_{k-1}^- = k \rangle \langle k $	$(k-1) \bullet \rightleftharpoons \bullet k \equiv \bigcirc k \equiv \bullet$

(4.8)

Using these rules, one sees that for the quintic the generic quiver diagram for the Z_i operators is given by a polygon with five vertices (a pentagon) and in general twenty links

joining the various vertices; see figure 1. Since a vector basis type $\mathbf{e}_{k(k+n)} = |k \rangle \langle k+n|$ can be usually decomposed as the product of n basis links as shown on this equation

$$\mathbf{e}_{k(k+n)} = \mathbf{a}_k^+ \mathbf{a}_{k+1}^+ \dots \mathbf{a}_{k+n}^+,$$

where $\mathbf{a}_k^+ = \mathbf{e}_{k(k+1)}$, one concludes that points in NC geometry are roughly speaking described by pentagons.




Figure 1: A pentagon diagram representing points of the NC quintic, $\mathcal{Q}^{nc}$. Since generic links may usually be decomposed in terms of the $|j \rangle \langle j+1|$ edges, this quiver diagram can also be thought as a polygon of five vertices and so NC points of $\mathcal{Q}^{nc}$ may be imagined as pentagons.

We will be use later on these diagrams to study the the fractional D brane at singularities of the orbifold of the quintic.

4.2 Discrete Torsion Matrix of $\mathbf{Z}_5^3$

Here we want to study general $\mathcal{D}(G)$ representations of the $\mathbf{Z}_5^3$ in presence of torsion. We shall first consider the simplest situation where all $\mathbf{Z}_5$ factors commute amongst themselves; then we discuss the case where they do not commute.

4.2.1 Free torsion case

Naively, the geometric $\mathbf{Z}_5^3$ can be seen as the product of three abelian $\mathbf{Z}_5$ group factors whose generators may be defined, by help of the tensor product, as follows:

$$\begin{aligned} E_1 &= P_1 \otimes I_{id} \otimes I_{id}, \\ E_2 &= I_{id} \otimes P_2 \otimes I_{id}, \\ E_3 &= I_{id} \otimes I_{id} \otimes P_3, \end{aligned} \tag{4.9}$$

The E'_a s are the generators of the three $\mathbf{Z}_5$ factors of $\mathbf{Z}_5^3$; they satisfy the cyclic property $E_a^5 = \mathbf{I}_{\mathcal{D}(G)}$ following from the individual identities $P_a^5 = I_{id}$. Since they are commuting operators; i.e,

$$E_a \cdot E_b = E_b \cdot E_a, \tag{4.10}$$

they can be diagonalized simultaneously in the same basis $\{|a, i\rangle; 1 \leq a \leq 3; 1 \leq i \leq 5\}$. As such the P'_a s can be thought more a less as in eq(3.20) and E'_a s as blocks of diagonal matrices. Using the convention notations $\mathbf{a}_{n_a}^+ = |a, n\rangle \otimes \langle a, n+1|$, $\mathbf{a}_{n_a}^- = |a, n+1\rangle \otimes \langle a, n|$ and $\pi_{n_a} = \mathbf{a}_{n_a}^+ \mathbf{a}_{n_a}^-$, and the graphic representation eqs(4.8), we can also represent the following operators decompositions;

$$\begin{aligned} I_{id} &= \sum_{n=1}^5 \pi_n, & P_a &= \sum_{n=1}^5 \alpha_{a,n} \pi_n, \\ Q_a &= \sum_{n=1}^5 \mathbf{a}_{a,n}^+; & Q_a^{-1} &= \sum_{n=1}^5 \mathbf{a}_{a,n}^-, \end{aligned} \tag{4.11}$$

in terms of quiver diagrams as shown on figures 2. For explicit computations, we will drop the index a by working in special matrix realizations.

Generic elements g of $\mathbf{Z}_5^3$ split also as $g_1 \otimes g_2 \otimes g_3$, the tensor product of the g_a elements of the $\mathbf{Z}'_5$ s. Since $\mathbf{Z}_5^3$ is a cyclic group, one can expand these the g' s, in terms of the E_a generators as follows:

$$g = \prod_{i_1, i_2, i_3=1}^5 \gamma_{i_1 i_2 i_3} E_1^{i_1} E_2^{i_2} E_3^{i_3}, \tag{4.12}$$




Figure 2: The completely reducible diagram Fig2.a represents the identity operator of the five dimensional representation $\mathcal{D}(\mathbf{Z}_5)$. Fig2.b is an oriented pentagon representing the Q automorphism operators while fig2.c its inverse.

where the $\gamma_{i_1 i_2 i_3}$ coefficients are such that $\gamma_{i_1 i_2 i_3}^5 = 1$. Indeed, since the group multiplication law on $\mathbf{Z}_5^3$ is defined as usual by performing multiplications of individual elements; that is $g.g' = g_1.g'_1 \otimes g_2.g'_2 \otimes g_3.g'_3$; we have $E_a^5 = \mathbf{I}_{\mathcal{D}(\mathbf{Z}_5^3)}$, with $\mathbf{I}_{\mathcal{D}(\mathbf{Z}_5^3)}$ standing for $I_{\mathcal{D}(\mathbf{Z}_5)} \otimes I_{\mathcal{D}(\mathbf{Z}_5)} \otimes I_{\mathcal{D}(\mathbf{Z}_5)} \equiv I_{id} \otimes I_{id} \otimes I_{id}$ group representation identity.

As all elements of $\mathbf{Z}_5^3$ can be expressed as powers of E_a 's, we will focus our attention on these generators as well as on their monomials $\{E_1^{n_1} E_2^{n_2} E_3^{n_3}; 1 \leq n_a \leq 5\}$. The latter generate the dimension $d_1 d_2 d_3$ group algebra $\mathcal{A}_G$ of $\mathbf{Z}_5^3$. The d_i integers are the dimensions of the three $\mathcal{D}(\mathbf{Z}_5)$ factors of $\mathcal{D}(\mathbf{Z}_5^3)$.

4.2.2 Discrete Torsion

Though abelian, the three $\mathbf{Z}_5$ factors of $\mathbf{Z}_5^3$ may also not commute among each others and this situation corresponds to the case where there exist discrete torsions. In other words given a couple of elements $(g_1, g_2) \in \mathbf{Z}_5 \otimes \mathbf{Z}_5$, then $g_1 g_2 \neq g_2 g_1$. In practice, we shall have $g_1 g_2 = \alpha g_2 g_1$ with α is a fifth root a unity. This situation, which is quite similar to the Fuzzy torii realizations known also as the rational non commutative torii, may be described by extending the relations (4.7) by allowing twisting co-cycles t_{ab} between the three $\mathbf{Z}_5^3$ factors as shown here below,

$$F_a F_b = t_{ab} F_b F_a, \quad (4.13)$$

where $t(F_a, F_b) \equiv t_{ab}$, is the torsion matrix and where the F'_a s are the new $\mathbf{Z}_5^3$ generators; they are essentially the E_a 's times some T_a matrices carrying the torsion effects; $F_a = T_a E_a$. Moreover, like the E_a 's and T_a 's, the F_a 's satisfy equally $F_a^5 = \mathbf{I}_{\mathcal{D}(\mathbf{Z}_5^3)}$. From the relations (4.10), one learns several informations on their representations:

Structure of the torsion matrix Using the cyclic group property $F_a^5 = \mathbf{I}_{\mathcal{D}(\mathbf{Z}_5^3)}$, one sees that the torsion matrix t_{ab} should be like $t_{ab} = \omega^{\theta_{ab}}$ where $\theta_{ab} = -\theta_{ba}$ is antisymmetric 3×3 matrix. For $\theta = 0$, one gets just the relations (4.7). A tricky parameterization of torsion is to take θ_{ab} as $\theta_{ab} = \eta_{ab} - \eta_{ba}$, where η_{ab} is an $SL(3, \mathbf{Z})$ matrix, and rewrite eqs(4.10) as

$$\omega^{\eta_{ba}} F_a F_b = \omega^{\eta_{ab}} F_b F_a. \quad (4.14)$$

Now using the change $F_a = T_a E_a$, one sees that the T_a operators are just the generators of automorphisms of $\mathbf{Z}_5^3$ as shown on the following relation

$$T_b E_a T_b^{-1} = \omega^{\eta_{ab}} E_a \quad (4.15)$$

Actually this relation should be compared with eqs(2.18) and (4.19-20). Therefore the T_a 's are just the generators of $\mathbf{Z}_5^{aut\ 3}$, the automorphism group of $\mathbf{Z}_5^3$ symmetry.

$\mathcal{D}(G)$ Representations Taking the determinant of eqs(4.14), one discovers that the dimension d of the $\mathcal{D}(G)$ matrix representations splits as the sum of three terms as given herebelow;

$$d = d_1 + d_2 + d_3 \in 5\mathbf{Z}, \quad (4.16)$$

where (d_1, d_2, d_3) are respectively the dimensions of the $\mathcal{D}(\mathbf{Z}_5) \otimes \mathcal{D}'(\mathbf{Z}_5) \otimes \mathcal{D}''(\mathbf{Z}_5)$ representations. From this decomposition, one sees that the F_a 's may be realized in various ways in terms of tensor products of $d_a \times d_a$ matrices. A convenient parameterization of the F_a generators, which we will use to establish our results, is given by the following;

$$\begin{aligned} F_1 &= P_1 \otimes Q_2^{\eta_{12}} \otimes Q_3^{\eta_{13}}, \\ F_2 &= Q_1^{\eta_{21}} \otimes P_2 \otimes Q_3^{\eta_{23}}, \\ F_3 &= Q_1^{\eta_{31}} \otimes Q_2^{\eta_{32}} \otimes P_3, \end{aligned} \quad (4.17)$$

where the P'_a s and the Q'_a s are as in eq(3.19) with $P_a Q_b = \alpha_a Q_b P_a \delta_{ab}$, and where the η_{ab} 's are integers. The latters are in fact some given integer numbers carrying torsion effect; they are the same as in eq(4.15). Viewed as a matrix, η_{ab} should be in $SL(3, \mathbf{Z})$; $\eta_{ab} \in SL(3, \mathbf{Z})$; and moreover should have a non vanishing antisymmetric part;

$$\begin{aligned} \eta_{ab} - \eta_{ba} &\neq 0, \quad \eta_{ab} \cdot \eta_{cb}^{-1} = \delta_{ac}, \\ \eta_{11}\eta_{22}\eta_{33} + \eta_{12}\eta_{13}\eta_{31} + \eta_{13}\eta_{11}\eta_{22} &\neq \eta_{11}\eta_{23}\eta_{32} + \eta_{12}\eta_{11}\eta_{33} + \eta_{13}\eta_{12}\eta_{21}. \end{aligned} \quad (4.18)$$

To fix the ideas, we set for simplicity $\alpha_1 = \alpha_2 = \alpha_3 = \omega$ and consider the special case where the three basis of the three representation factors of $\mathcal{D}(G)$, namely $\{|a, i\rangle; 1 \leq i \leq d_a; 1 \leq a \leq 3\}$, have the same dimension; i.e, $d_1 = d_2 = d_3$ and so $d = d_1 + d_2 + d_3 = 3d_1$. Extensions to general cases are straightforward and some of their aspects will be given in the conclusion. In this case and for $d_1 = 5$, the $\mathbf{P}_1(\omega) = \mathbf{P}_2(\omega) = \mathbf{P}_3(\omega) = \mathbf{P}$ and $\mathbf{Q}_1 = \mathbf{Q}_2 = \mathbf{Q}_3 = \mathbf{Q}$, are as in eq(3.19). With this choice, the relations $\mathbf{P}_a \mathbf{Q}_b = \alpha_a \mathbf{Q}_b \mathbf{P}_a \delta_{ab}$ reduce to tree identical equations of type;

$$\begin{aligned} \mathbf{P} \cdot \mathbf{Q} &= \omega \mathbf{Q} \cdot \mathbf{P}, \\ \mathbf{P}^5 &= I_{id}, \quad \mathbf{Q}^5 = I_{id}. \end{aligned} \quad (4.19)$$

Before going ahead, we want to give three remarks and make some comments regarding eqs(4.17) and (4.18-19). The remarks regard the parameterization (4.17), the automorphism symmetry and the $\mathcal{D}(G)$ representations.

The F_a Basis The parameterization eq(4.17) is very special, first because it is related to the abelian one as

$$F_a = T_a E_a, \quad (4.20)$$

where the T_a 's are operators built out of monomials of the following generators;

$$\begin{aligned} J_1 &= Q_1 \otimes I_{id} \otimes I_{id}, \\ J_2 &= I_{id} \otimes Q_2 \otimes I_{id}, \\ J_3 &= I_{id} \otimes I_{id} \otimes Q_3, \\ J_a^5 &= \mathbf{I}_{\mathcal{D}(G)}. \end{aligned} \quad (4.21)$$

Indeed, comparing eq(4.20) with eqs(4.17), it is not difficult to check that the T_a 's read as follows;

$$\begin{aligned}
T_1 &= J_2^{\eta_{12}} \cdot J_3^{\eta_{13}}, \\
T_2 &= J_1^{\eta_{21}} \cdot J_3^{\eta_{23}}, \\
T_3 &= J_1^{\eta_{31}} \cdot J_2^{\eta_{32}}, \\
T_a^5 &= \mathbf{I}_{\mathcal{D}(G)}.
\end{aligned} \tag{4.22}$$

This means that torsion will be generated by the J_a operators. The second thing we want to mention is that there are more general parameterizations F'_a for the group generators going beyond eqs(4.17). They extend the F_a 's and involve E_a 's and J_a 's on equal footing as well as two kind of torsion matrices n_{ab} and ν_{ab} as shown here below;

$$\begin{aligned}
F'_1 &= P_1^{n_{11}} Q_1^{\nu_{11}} \otimes P_2^{n_{12}} Q_2^{\nu_{12}} \otimes P_3^{n_{13}} Q_3^{\nu_{13}}, \\
F'_2 &= P_1^{n_{21}} Q_1^{\nu_{21}} \otimes P_2^{n_{22}} Q_2^{\nu_{22}} \otimes P_3^{n_{23}} Q_3^{\nu_{23}}, \\
F'_3 &= P_1^{n_{31}} Q_1^{\nu_{31}} \otimes P_2^{n_{32}} Q_2^{\nu_{32}} \otimes P_3^{n_{33}} Q_3^{\nu_{33}},
\end{aligned} \tag{4.23}$$

or equivalently in terms of powers of the E_a 's and J_a 's;

$$F'_a = \prod_{b=1}^3 (E_1^{n_{ab}} J_1^{\nu_{ab}} E_2^{n_{ab}} J_2^{\nu_{ab}} E_3^{n_{ab}} Q_3^{\nu_{ab}}), \tag{4.24}$$

where n_{ab} and ν_{ab} are integers modulo 5. Eqs(4.23) can be related as well to the abelian one (4.10); but also to the parameterization eqs(4.17) as

$$F'_a = T'_a F_a, \tag{4.25}$$

where T'_a 's are now monomials of the primitive E_a and J_a generators; i.e $T_a = T_a(E_1, E_2, E_3; J_1, J_2, J_3)$ with $T_a^5 = \mathbf{I}_{\mathcal{D}(G)}$. Due to this arbitrariness in the choice of the basis of generators of the $\mathcal{D}(G)$ group representation, one learns already at this level that there are different ways to realize the non commutative extension of the quintic. The so called Z and Φ realizations described in section 2 are just two special ones.

More on Quantum Symmetries The J_a operators have a nice interpretation; they are the primitive generators of the $\mathbf{Z}_5^{aut \otimes 3}$ automorphism group of the $\mathbf{Z}_5^3$ symmetry of the quintic. This is clearly seen on eqs(4.24), which associate to each symmetry generator E_a , an automorphism J_a . Eqs(4.23-24) tell us that E_a and J_a generate three copies of non commutative two dimensional fuzzy torii of parameters α_a ; $\mathcal{T}_{\alpha_1} \otimes \mathcal{T}_{\alpha_2} \otimes \mathcal{T}_{\alpha_3}$; and moreover show that the full symmetry of the quintic is in fact $(\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3$. The latter acts on the z complex variables as as in eqs(3.2) and (3.15)

Furthermore, the analysis we have been developing here for discrete torsion is valid for any complex d dimension Calabi-Yau orbifolds $\mathcal{O}_d$ with discrete group $\mathbf{Z}_{d+2}^{\otimes d}$ generating a real $2d$ dimensional NC torus of type $\mathcal{T}_{\alpha_1} \otimes \dots \otimes \mathcal{T}_{\alpha_d}$. It is also suspected to apply as well to the continuous limit $\mathbf{Z}_\infty \simeq U(1)$ where we expect the appearance of irrational representations of non commutative Torii and Powers-Riffel projectors [25-28] together with a continuous fractional D -brane spectrum. Further details on the last issue will be reported in [33]

Representations The constraint eq(4.16) on the dimensions of the F_a representations shows that it is possible to build different realizations for the F_a generators. The representation eqs(3.19) built in [24], corresponds to take (d_1, d_2, d_3) equal to either $(5, 0, 0)$, $(0, 5, 0)$, or $(0, 0, 5)$ respectively obtained by setting $\eta_{a2} = \eta_{a3} = 0$, $\eta_{a1} = \eta_{a3} = 0$ and $\eta_{a1} = \eta_{a2} = 0$. Here we shall give new solutions of type (d_1, d_2, d_3) by keeping all $\eta_{ab} \neq 0$.

Now, using the relations(4.17), one can compute the expression of the t_{ab} cocycles between the F_a generators. We find

$$t(F_a, F_b) = \omega^{(\eta_{ab} - \eta_{ba})}, \quad (4.26)$$

showing that θ_{ab} is equal to $(\eta_{ab} - \eta_{ba})$. As we will handle much more θ_{ab} and η_{ab} than the $t(F_a, F_b)$ matrix, we will also refer to θ_{ab} and η_{ab} as discrete torsion matrices of the group $\mathbf{Z}_5^3$; though they are proportional to its logarithm; i.e $\theta_{ab} = \frac{-5i}{1+5k} \text{Log}(t_{ab})$; $k \in \mathbf{Z}$. Note in passing that torsion is carried by the antisymmetric part of the matrix η_{ab} . We will also see that the antisymmetric part of its inverse $\eta_{[ab]}^{-1}$ plays equally a crucial role in working out the solutions for the NC quintic.

4.2.3 Quiver Diagrams

Like for the case of one abelian factor, one can also build the projectors for full the $\mathbf{Z}_5^3$ group. Using the individual $\mathbf{Z}_5$ projectors $\pi_{k_a} = \frac{1}{5} \sum_{i=1}^5 \omega^{(-k_a)} \mathbf{P}_\omega^i$, we can construct a

variety of projectors on the representation space of $\mathbf{Z}_5^3$. First, the Π_{k_a} projectors on the $\mathbf{Z}_5$ representation spaces;

$$\begin{aligned}\Pi_{k_1} &= \pi_{k_1} \otimes I_{id} \otimes I_{id}, \\ \Pi_{k_2} &= I_{id} \otimes \pi_{k_2} \otimes I_{id}, \\ \Pi_{k_3} &= I_{id} \otimes I_{id} \otimes \pi_{k_3},\end{aligned}\tag{4.27}$$

They have quiver diagrams more or less similar to that of π_{k_a} , except that now we have some richness coming from the decomposition of the extra identity operators. In any way, the full quiver diagram is given by the cross product of the individual graphs.

Second, the $\Pi_{(k_a, k_b)}$ and $\Pi_{(k_1, k_2, k_3)}$ projectors on the $\mathbf{Z}_5^2$ and $\mathbf{Z}_5^3$ representation spaces, respectively obtained by taking products of Π'_{k_a} s;

$$\begin{aligned}\Pi_{(k_a, k_b)} &= \Pi_{k_a} \Pi_{k_b} \\ \Pi_{(k_1, k_2, k_3)} &= \Pi_{k_1} \Pi_{k_2} \Pi_{k_3}.\end{aligned}\tag{4.28}$$

Accordingly, the identity matrix $\mathbf{I}_{\mathcal{D}(G)}$ can be decomposed in different ways as shown herebelow;

$$\mathbf{I}_{\mathcal{D}(G)} = \sum_{k_a=1}^5 \Pi_{k_a} = \sum_{k_a, k_b=1}^5 \Pi_{k_a, k_b} = \sum_{k_1, k_2, k_3=1}^5 \Pi_{k_1} \Pi_{k_2} \Pi_{k_3}.\tag{4.29}$$

So $\mathbf{I}_{\mathcal{D}}$ can be represented by a completely reducible Quiver diagram with $d_1 d_2 d_3 = 5 \times 5 \times 5$ vertices. Similar expansion to eqs(4.29) may be written down for the generators J_a of the quantum symmetries. Setting

$$\begin{aligned}A_{k_1}^{\pm} &= a_{1, k_1}^{\pm} \otimes I_{id} \otimes I_{id}, \\ A_{k_2}^{\pm} &= I_{id} \otimes a_{1, k_1}^{\pm} \otimes I_{id}, \\ A_{k_3}^{\pm} &= I_{id} \otimes I_{id} \otimes a_{k_3}^{\pm},\end{aligned}\tag{4.30}$$

we can write for instance J_a and $J_{a_1} J_{a_2}$ as follows;

$$J_a = \sum_{k_a=1}^5 A_{k_a}^+, \quad J_{a_1} J_{a_2} = \sum_{k_{a_1} k_{a_2}=1}^5 A_{k_{a_1}}^+ A_{k_{a_2}}^+.\tag{4.31}$$

While the quiver diagram for the J_a are similar to that given by figures 2.b and c and figures 3, the quiver diagrams associated with $J_{a_i} J_{a_j}$ are obtained by taking cross products and are of type figure 4.




Figure 3: Fig 3-a represents the quiver diagram of the Q^2 operator, the oriented links define the independent massless chiral fields on the D brane at singularity. Fig 3-b represents the quiver diagram of the Q^3 , which upon reversing the orientation of the links is similar to fig 3-a. Fig 3-c represents the quiver diagram of the Q^4 operator, it just the reverse of the diagram of Q and Fig 3-d represents the completely reducible quiver diagram of the Q^5 .

5 CP Algebra

The construction we will develop here applies to a wide class Calabi-Yau orbifolds with torsion and offer a unified description for the study of non commutative geometry versus discrete torsion. It explains the origin of the various possible realizations of non commutative manifolds encountered in the previous section; see eqs(4.17) and (4.23). The strategy we will be following in this presentation is: First build the CP algebra for the quintic by working in the Φ_i frame. We study its centre $\mathcal{Z}(\mathcal{A}[\Phi, F])$ and the structure of the non commutative points of $\mathcal{Q}$. Then, we build a class of explicit solutions for the fields moduli by taking Φ_i 's as matrix operators valued in $\mathcal{D}(\mathbf{Z}_5^{aut^3})$. Next we reconsider the Z realization from the point of the CP algebra method and then build the Morita equivalence between $\mathcal{A}[Z, F]$ and $\mathcal{A}[\Phi, F]$. Finally we consider other frames of the CP algebra and their Morita equivalence.

5.1 $Q \boxtimes \mathbf{Z}_5^3$ CP Algebra

Roughly speaking the CP algebra of the quintic $Q \boxtimes \mathcal{A}_G$, with $G = \mathbf{Z}_5^3$, is a subspace of the algebra of functionals $\mathcal{F}[[\mathcal{M}at(d_1 d_2 d_3, \mathbf{C})]]$ on the tensor product of the $\mathcal{M}at(d_a, C)$ matrix spaces. This is a fiber bundle based on Q and whose fiber is given by the algebra of the representation group $\mathcal{D}(G)$. Elements of $Q \boxtimes \mathcal{A}_G$ are generated by the matrix fields moduli Φ_i , $1 \leq i \leq 5$, and the $\mathcal{D}(G)$ group generators F_a , $1 \leq a \leq 3$. The Φ_i chiral operators and the

F_a generators satisfy the following algebra; see also eqs(2.18) and (2.20),

$$\begin{aligned}\Phi_i \cdot \Phi_j &= \Phi_j \cdot \Phi_i, \\ F_a \cdot \Phi_i &= \lambda_{ai} \Phi_i \cdot F_a, \\ F_a \cdot F_b &= \omega^{\theta_{ab}} F_b \cdot F_a,\end{aligned}\tag{5.1}$$

where λ_{ai} can be set equal to $\omega^{q_i^a}$ as required by the constraint eq $F_a^5 = \mathbf{I}_{\mathcal{D}(G)}$. The q_i^a 's are the same integers we have been considering so far and, for simplicity, will be chosen as in eqs(3.4). The θ_{ab} 's are integers defining the discrete torsion matrix eqs(4.11). They are fixed numbers for well specified torsions. Actually the relations (5.1) define the Φ realization of the NC quintic $\mathcal{Q}^{(nc)}$. A class of solutions for $\mathcal{Q}^{(nc)}$ were already considered in section 3 by using the constrained method. In addition to its direct physical interpretation as supersymmetric vacuum configurations and its link with $4d$ superpotentials (2.13), there are other novelties in the Φ realization are: (1) The matrix fields moduli Φ_i are commuting operators contrary to the Z_i 's which do not as shown on eqs(3.7). (2) The number of generators in the CP algebra of the quintic is apparently larger than those appearing in eqs(3.7). In addition to the Φ_i 's, we have moreover the group generators as principal actors. In eqs(3.7), the F_a group generators act as discrete outer automorphisms; but in the Φ realization they are inner ones.

To prove that the algebra generated by eqs(5.1) is indeed the non commutative extension of the quintic eq(3.1), we have to show that: (i) the centre $\mathcal{Z}(Q \boxtimes \mathcal{A}_G)$ of the CP algebra eq(5.1) is exactly the commutative quintic $\mathcal{Q}$ given by eq(3.1) and (ii) check that eqs(5.1) are equivalent to eqs(2.22) and (3.9) used for defining $\mathcal{Q}^{(nc)}$ in the Z realization. Put differently, we have to establish that,

$$\begin{aligned}Q \boxtimes \mathcal{A}_G &\simeq \mathcal{Q}^{(nc)}, & (a) \\ \mathcal{Z}(Q \boxtimes \mathcal{A}_G) &\simeq \mathcal{Q}. & (b)\end{aligned}\tag{5.2}$$

We will first address eq(5.2-a) and then discuss the equivalence.

5.1.1 The Centre $\mathcal{Z}\mathcal{A}(Q \boxtimes \mathcal{A}_G)$

To start note that generic elements v of $Q \boxtimes \mathcal{A}_G$ are functions of the matrix generators Φ_i and F_a ; i.e, $v = v(\Phi_1, \dots, \Phi_5; F_1, F_2, F_3)$. As the $\mathbf{Z}_5$'s are cyclic groups, the v 's can be usually expanded in a power series of the $\mathcal{D}(G)$ generators as follows;

$$v = \sum_{i,j,k=1}^5 \mathcal{F}_{ijk}(\Phi_1, \dots, \Phi_5) F_1^i F_2^j F_3^k, \quad (5.3)$$

where $\mathcal{F}_{ijk}(\Phi_1, \dots, \Phi_5)$ are arbitrary analytic functions of the Φ_l matrix fields. To get the elements $\zeta_\mu = \zeta_\mu(\Phi_1, \dots, \Phi_5)$ of the centre $\mathcal{Z}(Q \boxtimes \mathcal{A}_G)$, one has to solve the constraint eqs

$$\zeta_\mu v = v \zeta_\mu, \quad \forall v \in Q \boxtimes \mathcal{A}_G, \quad (5.4)$$

or equivalently,

$$\begin{aligned} \zeta_\mu \Phi_i &= \Phi_i \zeta_\mu, \\ \zeta_\mu F_a &= F_a \zeta_\mu. \end{aligned} \quad (5.5)$$

Eq(5.4) shows that, in presence of discrete torsion, $\zeta_\mu = \mathcal{F}_{(0,0,0)}(\Phi_1, \dots, \Phi_5)$ while eqs(5.5) requires that ζ_μ should be invariant under the $\mathbf{Z}_5^3$ actions. Particular solutions of these constraint eqs, in agreement with the Schur lemma, are given by,

$$\begin{aligned} \zeta_i &= \Phi_i^5 = x_i^5 \mathbf{I}_{\mathcal{D}(G)} \\ \zeta_0 &= \Pi_{i=1}^5 \Phi_i = \mathbf{I}_{\mathcal{D}(G)} (\Pi_{i=1}^5 x_i), \end{aligned} \quad (5.6)$$

where the x_i 's are given by (2.3); they are essentially as in eq(3.1). Therefore the centre $\mathcal{Z}(Q \boxtimes \mathcal{A}_G)$ is given by the linear combination of ζ_μ , namely $\sum_\mu \zeta_\mu = \sum_{i=1}^5 (x_i^5 + \Pi_{i=1}^5 x_i) \mathbf{I}_{\mathcal{D}(G)}$. As such, generic v elements of the quintic CP algebra can be usually rewritten as:

$$v = \sum_{i=1}^5 (x_i^5 + \Pi_{i=1}^5 x_i) \mathbf{I}_{\mathcal{D}(G)} + \sum_{i,j,k=1}^5 \mathcal{F}'_{k_1 k_2 k_3}(\Phi_1, \dots, \Phi_5) F_1^{k_1} F_2^{k_2} F_3^{k_3}. \quad (5.7)$$

Actually this relation shows that non commutative points in the $(Q \boxtimes \mathcal{A}_G)$ CP algebra of the quintic can be thought of as fibers of a fiber bundle of base $\sum_{i=1}^5 (x_i^5 + \Pi_{i=1}^5 x_i) = 0$ and fiber given by the algebra of the group representation $\mathcal{D}(G)$.

5.1.2 Structure of the Fibration

Demanding invariance of the above relation under $\mathbf{Z}_5^3$, we can extract some informations on the structure of the $\mathcal{F}'_{k_1 k_2 k_3}$ monomials. Indeed, solving the identity $F_a v F_a^{-1} = v$, we get;

$$F_a \mathcal{F}'_{k_1 k_2 k_3} F_a^{-1} = Q_{(k_1 k_2 k_3)}^a \mathcal{F}'_{k_1 k_2 k_3}, \quad (5.8)$$

with

$$Q_{(k_1 k_2 k_3)}^a = - \sum_{b=1}^3 k_b \theta_{ab} + 5\mathbf{Z}. \quad (5.9)$$

Moreover, using the relation $\eta_{ab} l_{ij} = q_i^a q_j^b$ that follows from eqs(3.14-15), one can express θ_{ab} as

$$\theta_{ab} = \sum_{i,j=1}^5 q_i^a (l_{ij}^{-1} - l_{ji}^{-1}) q_j^b \quad (5.10)$$

Putting back into eq(5.9), we find

$$Q_{(k_1 k_2 k_3)}^a = - \sum_{i=1}^5 q_i^a \left[\sum_{j=1}^5 \sum_{b=1}^3 k_b (l_{ij}^{-1} - l_{ji}^{-1}) q_j^b \right] + 5\mathbf{Z}. \quad (5.11)$$

But this relation tells us that the $\mathcal{F}'_{k_1 k_2 k_3}$ monomials have the form

$$\mathcal{F}'_{k_1 k_2 k_3} = \prod_{i,j=1}^5 \left(\left[(\Phi_i)^{(l_{ij}^{-1} - l_{ji}^{-1}) q_j^1} \right]^{k_1} \left[(\Phi_i)^{(l_{ij}^{-1} - l_{ji}^{-1}) q_j^2} \right]^{k_2} \left[(\Phi_i)^{(l_{ij}^{-1} - l_{ji}^{-1}) q_j^3} \right]^{k_3} \right) \quad (5.12)$$

Therefore the most general form of the non commutative quintic invariant under $\mathbf{Z}_5^3$ transformation is given by eqs(5.3) with $\mathcal{F}'_{k_1 k_2 k_3}$ as in eqs(5.12).

5.2 Explicit Expression of the Φ_i 's

Explicit solutions for the generators of the $Q \boxtimes \mathcal{A}_G$ algebra can be written down by using the commutativity feature of the Φ_i 's. To build a matrix representation, we consider the simple choice eqs(4.17) for the F_a group generators;

$$\begin{aligned} F_1 &= P \otimes Q^{\eta_{12}} \otimes Q^{\eta_{13}}, \\ F_2 &= Q^{\eta_{21}} \otimes P \otimes Q^{\eta_{23}}, \\ F_3 &= Q^{\eta_{31}} \otimes Q^{\eta_{32}} \otimes P, \end{aligned} \quad (5.13)$$

where $P_1 = P_2 = P_3 = P$ and where P and Q satisfy eqs(3.20) with α taken as the fifth primitive root of unity. The F_1, F_2 and F_3 operators are mainly elements of $\mathcal{D}(\mathbf{Z}_5 \otimes \mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5^{aut})$, $\mathcal{D}(\mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5 \otimes \mathbf{Z}_5^{aut})$ and $\mathcal{D}(\mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5)$ respectively. Using these relations, it is not a difficult to check that,

$$F_1 \left[Q^{q_1} \otimes Q^{q_2} \otimes Q^{q_3} \right] = \omega^{q_1} \left[Q^{q_1} \otimes Q^{q_2} \otimes Q^{q_3} \right] F_1. \quad (5.14)$$

More generally we have the following identities;

$$F_a \left[J_1^{q_1} . J_2^{q_2} . J_3^{q_3} \right] = \omega^{q_a} \left[J_1^{q_1} . J_2^{q_2} . J_3^{q_3} \right] F_a. \quad (5.15)$$

Therefore a class of solution for the matrix fields moduli reads as

$$\begin{aligned} \Phi_i &= z_i \left[Q^{q_1} \otimes Q^{q_2} \otimes Q^{q_3} \right] \\ &= z_i \prod_{a=1}^3 J_a^{q_a}. \end{aligned} \quad (5.16)$$

Such class of solutions belongs to the algebra of the representation of the $\mathcal{D}(\mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5^{aut} \otimes \mathbf{Z}_5^{aut})$ representation of the automorphism symmetry. In addition to the manifest commutativity property $\Phi_i . \Phi_j = \Phi_j . \Phi_i$, it satisfies naturally the relations (5.1-2); due to the property $Q^5 = I_{id}$ and to the Calabi-Yau condition eqs(3.5). From these solutions one also learns that it is possible to build other solutions by using both $\mathbf{Z}_5$ and $\mathbf{Z}_5^{aut}$ factors of the full discrete symmetry $\widehat{G}$. In what follows we will consider these kind of solutions and show that they are indeed equivalent to eqs(5.13) and (5.16).

6 General Solutions

Let us summarize what we have been studying so far. The algebraic relations defining the non commutative quintic $\mathcal{Q}^{(nc)}$ as appeared first in [2] read, in the coordinate patch $z_5 = 1$, as;

$$Z_1 Z_2 = \alpha Z_2 Z_1,$$

$$\begin{aligned}
Z_1 Z_3 &= \alpha^{-1} \beta Z_3 Z_1, \\
Z_1 Z_4 &= \beta^{-1} Z_4 Z_1, \\
Z_2 Z_3 &= \alpha \gamma Z_3 Z_2, \\
Z_2 Z_4 &= \gamma^{-1} Z_4 Z_2, \\
Z_3 Z_4 &= \beta \gamma Z_4 Z_3, \\
Z_i Z_5 &= Z_5 Z_i, \quad i = 1, 2, 3, 4;
\end{aligned} \tag{6.1}$$

where α, β, γ are fifth roots of unity. In [24], it was noted that the above relations are very special and can be generalized as follows:

$$\begin{aligned}
Z_i Z_j &= \beta_{ij} Z_j Z_i; \quad i, j = 1, \dots, (n+1), \\
Z_i Z_{d+2} &= Z_{d+2} Z_i; \quad i = 1, \dots, (n+1),
\end{aligned} \tag{6.2}$$

For the case of the quintic the n parameter should be set equal to three; $n = 3$; but the above relations are also valid for complex n -dimension holomorphic homogeneous Calabi-Yau hypersurfaces. The non commutative deformation parameters β_{kl} are realized in terms of the Calabi-Yau charges as follows:

$$\beta_{ij} = \exp i \left(\frac{2\pi}{n+2} m_{ab} q_i^a q_j^b \right) = \omega^{m_{ab} q_i^a q_j^b}. \tag{6.3}$$

where m_{ab} is some given matrix with integer entries; it has remained without interpretation in [24]; but here we will identify its origin. Indeed, using the CP algebra representation of the non commutative quintic, we will prove that m_{ab} is equal to $(\eta_{ab}^{-1} - \eta_{ba}^{-1})$; where η_{ab} is as in eqs(5.13). To that purpose, let us analyze the conditions under which eqs(5.1) and (6.1-2) are equivalent.

6.1 Morphisms between Eqs(5.1) and Eqs(6.1)

At first sight, saying that eqs (5.1) and eqs(6.1-2) have to be equivalent seems a little bit ambiguous as the naive counting of the degrees of freedom seems not matching. Eqs (6.1-2) associated with the constrained formulation of the non commutative quintic, involve only the space variable operators Z_i ; the F_a group generators do not appear manifestly; they act as outer automorphisms on eqs (6.1-2). This is however not a major problem since it can be

overcome if one supposes that in the constrained form, the Z_i 's and the F_a 's do not couple each others and so eqs(6.2) should be rather understood as:

$$\begin{aligned} Z_i Z_j &= \beta_{ij} Z_j Z_i; & i, j &= 1, \dots, (d+1), \\ Z_i Z_{d+2} &= Z_{d+2} Z_i; & i &= 1, \dots, (d+1), \end{aligned} \quad (6.4)$$

together with

$$F_a F_b = \omega^{\theta_{ab}} F_b F_a, \quad (6.5)$$

and moreover,

$$F_a Z_i = Z_i F_a. \quad (6.6)$$

Though trivial, eqs(6.6) are however very important for us; they will determine the morphism linking eqs(5.1) and eqs(6.1-2). To do so, let us first introduce, Ψ_L and Ψ_R , two maps describing respectively left and right morphisms of the algebra eqs(5.1):

$$\Psi_L : \Phi_i \longrightarrow Z_i = \Gamma_i \cdot \Phi_i, \quad (6.7)$$

$$\Psi_R : \Phi_i \longrightarrow Z'_i = \Phi_i \cdot \Gamma_i, \quad (6.8)$$

where Γ_i is an invertible operator depending on the generators of the algebra; $\Gamma_i = \Gamma_i(\Phi_1, \dots, \Phi_5; F_a)$. Here below, we will take the set of Γ'_i s in the group representation $\mathcal{D}(G)$ as,

$$\begin{aligned} \Gamma_i &= [P^{p_i^1} \otimes P^{p_i^2} \otimes P^{p_i^3}]; \\ &= E_1^{p_i^1} \cdot E_2^{p_i^2} \cdot E_3^{p_i^3}, \end{aligned} \quad (6.9)$$

where the p_i^a 's are for the moment some given integers modulo five; they will be determined by requiring the relations eqs(6.4) to be hold. Now putting eqs(6.9) back into eqs(6.7-8) and using the realization of Φ_i given by eq(5.16) as well as the eqs(5.14-15), one gets amongst others;

$$Z'_i = \omega^{-\sum_{a=1}^3 p_i^a q_i^a} Z_i. \quad (6.10)$$

Actually this relation shows that Ψ_L and Ψ_R are related; $\Psi_R = (\omega^{-\sum_{a=1}^3 p_i^a q_i^a}) \Psi_L$; so we will focus our attention here after on the left morphism Ψ_L only.

Next, we solve $F_a.Z_i = Z_i.F_a$ by using eqs(5.13) and eqs(6.7). We find:

$$\sum_{b=1}^3 \eta_{ab}.p_i^b = q_i^a, \quad (6.11)$$

where η_{ab} is the same matrix we have been using. This relation shows that the p_i^a are related to the q_i^a via the torsion matrix namely $p_i^a = \sum_{b=1}^3 \eta_{ab}^{-1} q_i^b$. Since the p_i^a are integers, eq(6.11) requires that the matrix η_{ab} to belong to $Sl(3, \mathbf{Z})$ and shows moreover that p_i^a 's satisfy themselves the Calabi-Yau condition:

$$\sum_{i=1}^5 p_i^a = 0. \quad (6.12)$$

Furthermore eqs(6.11) show that, whenever torsions are present, the commutative quintic admits an extra hidden discrete symmetry acting on the z_i 's as

$$z_i \rightarrow \omega^{p_i^a} z_i. \quad (6.13)$$

Therefore the Ψ_L morphism is torsion dependent and is given by the left multiplication operators;

$$\Gamma_i = \prod_{a=1}^3 (P_{\mathbf{a}}^{-\sum_{b=1}^3 \eta_{ab}^{-1} q_i^b}). \quad (6.14)$$

This is a completely reducible operator suggesting that the Ψ_L morphism can be also written as $\Psi_L = \Psi_1 \otimes \Psi_2 \otimes \Psi_3$, with $\Psi_a = \oplus_{i=1}^5 \Psi_{a,i}$ and $\Psi_{a,s} = \alpha^{-s \sum_{b=1}^3 \eta_{ab}^{-1} q_i^b}$. The Ψ_L morphism depends on the Calabi-Yau charges q_i^a and the inverse of the discrete torsion matrix η_{ab}^{-1} . Using these expressions of and combining altogether the above relations, we get the Z_i realization for the CP algebra of the non commutative quintic, namely

$$\begin{aligned} Z_i &= P^{p_i^1} Q^{q_i^1} \otimes P^{p_i^2} Q^{q_i^2} \otimes P^{p_i^3} Q^{q_i^2}, \\ &= \prod_{a=1}^3 E_a^{p_i^a} J_a^{q_i^a}, \end{aligned} \quad (6.15)$$

where p_i^a are as in eqs(6.11-12). Finally we calculate the relation between $Z_i Z_j$ and $Z_j Z_i$. Using eqs(5.16) and eqs(6.7-9), we show, by straightforward computations, that

$$Z_i Z_j = \omega^{\sum_{a=1}^3 [p_i^a q_j^a - p_j^a q_i^a]} Z_j Z_i \quad (6.16)$$

Comparing this relation with eqs(6.3), one gets $m_{ab} q_i^a q_j^b = [p_i^a q_j^a - p_j^a q_i^a]$, then using the inverse eq(6.11) by expressing p_i^a in terms of the q_i^a , we can rewrite the previous relation as $m_{ab} q_i^a q_j^b = (\eta_{ba}^{-1} - \eta_{ab}^{-1}) q_i^a q_j^b$ showing that $m_{ab} = -(\eta_{ab}^{-1} - \eta_{ba}^{-1})$. In what follows we collect the various relations one gets,

$$\begin{aligned} l_{ij} &= \sum_{a=1}^3 p_i^a q_j^a = \sum_{a,b=1}^3 \eta_{ab}^{-1} q_i^b q_j^a = \sum_{a=1}^3 \eta_{ab} p_i^a p_j^b \\ L_{ij} &= l_{ij} - l_{ji} = p_i^a q_j^a - p_j^a q_i^a, \\ L_{ij} &= (\eta_{ab} - \eta_{ba}) p_i^a p_j^b, \\ L_{ij} &= -(\eta_{ab}^{-1} - \eta_{ba}^{-1}) q_i^a q_j^b. \end{aligned} \quad (6.17)$$

We end this discussion on the non commutative quintic representations by noting that given: (i) a set of Calabi-Yau charges q_i^a , defining the charges of $\mathbf{Z}_5^3$, and (ii) an invertible matrix η_{ab} of $SL(3; \mathbb{Z})$ with a non zero antisymmetric part; we can build various, but equivalent, realizations of the CP algebra of the NC quintic. Starting from eqs(5.16), perform variables change type;

$$\begin{aligned} \Phi'_i &= \Omega_i \Phi_i, \\ \Phi'_i &= \tau_i \Phi_i \Omega_i, \\ \Omega_i \Omega_j &= \varepsilon_{ij} \Omega_j \Omega_i, \\ F_a \Omega_i &= \kappa_{ai}^{-1} \Omega_i F_a, \end{aligned} \quad (6.18)$$

where $\Omega_i = \Omega_i(E_a, J_a)$ are given monomials of E_a and J_a with structure constants τ_i, ε_{ij} and κ_{ai} . Then compute the commutation relations for the new operators, we get;

$$\begin{aligned} \Phi'_i \Phi'_j &= \tau_i^{-1} \varepsilon_{ij} \tau_j \Phi'_j \Phi'_i, \\ F_a \Phi'_i &= \kappa_{ai}^{-1} \lambda_{ai} \Phi'_i F_a, \\ F_a F_b &= t_{ab} F_b F_a, \end{aligned} \quad (6.19)$$

where the new generators Φ'_i do no longer commute among themselves nor with the group representation generators. These relations are realizations of the NC quintic interpolating between the Φ_i realization and the Z_i one.

6.2 Example

To fix the ideas, let consider an example by choosing Ω_i as $F_1^{r_1^a} F_2^{r_2^a} F_3^{r_3^a}$ with r_i^a some integers. In this case, the above mentioned structure constants read as;

$$\begin{aligned}\tau_i &= \omega^{\sum_a r_i^a q_i^a}, \\ \varepsilon_{ij} &= \omega^{\theta_{ab} r_i^a r_j^b}, \\ \kappa_{ai} &= \omega^{-\theta_{ab} r_i^b},\end{aligned}\tag{6.20}$$

As we see, the structure constants ε_{ij} and κ_{ai} are torsion dependent; if $\theta_{ab} = 0$; then $\varepsilon_{ij} = 1$ and $\kappa_{ai} = 1$. Note that in the $\{\Phi'_i; F_a\}$ frame, the generators of the CP algebra of the quintic do not commute in general; except for the following special cases where the non commutative quintic takes remarkable forms:

(1) Φ_i Frame: eqs(5.1) and eqs(5.16)

This is recovered by setting $\tau_i^{-1} \varepsilon_{ij} \tau_j = 1$; that is,

$$\varepsilon_{ij} = \tau_i \tau_j^{-1}, \quad \kappa_{ai} = 1.\tag{6.21}$$

From eq(6.14), one sees that Ω_i is just the inverse of Γ_i ; that is, $\Omega_i = \Gamma_i^{-1}$.

(2) Z_i Frame: eqs(6.1-2) and eqs(6.15)

It is obtained by requiring the Z_i and F_a operators to commute; this is equivalent to setting $\kappa_{ai} \sigma_{ai} = 1$;

$$\begin{aligned}\sigma_{ai} &= \kappa_{ai}^{-1}, \quad \beta_{ij} = \tau_i^{-1} \varepsilon_{ij} \tau_j, \\ \Omega_i &= \Gamma_i.\end{aligned}\tag{6.22}$$

Since the two sets of matrix generators $\{\Phi'_i = Z_i\}$ and $\{F_a\}$ decouple completely, the NC quintic is then described by a trivial fibration of the group algebra $\mathcal{A}_G$ on the commutative quintic as shown here below;

$$Q \boxtimes \mathcal{A}_G \equiv \mathcal{A}[\Phi_i ; F_a] \equiv \mathcal{A}[Z_i] \otimes \mathcal{A}[F_a] \equiv \mathcal{Q}^{nc} \otimes \mathcal{A}_G. \quad (6.23)$$

This relation shows that the constrained method we studied in section 3 and the CP algebra approach we have been considering in sections 5 and 6 are Morita equivalent.

7 Fractional Branes

The realization of the NC quintic we have studied here above concerns only the regular points of the algebra, that is non singular ones. In this section, we want to complete this analysis by considering the representations for singular points. This is not only important for the study of fractional branes at singularities but also for answering the question regarding the nature of fractional branes on the NC quintic and more generally on NC Calabi-Yau hypersurfaces.

To do so, we shall first determine and classify the various sets $\mathcal{S}_{(\mu)}$ of singular points of orbifolds of the quintic; then we give the corresponding singular solutions. At first sight and as far as the full $\mathbf{Z}_5^3$ geometric symmetry of $\mathcal{Q}$ is concerned, we have only one fixed point under the $\mathbf{Z}_5^3$ actions, namely $(z_1, z_2, z_3, z_4, 1) = (0, 0, 0, 0, 1)$. This point belongs however to the $\mathbf{CP}^4$ projective space; but does not belong to the quintic $\mathcal{Q}$; no point of the quintic is then fixed by the full symmetry. This property is valid for all Calabi-Yau hypersurfaces; no point of complex n dimensional Calabi-Yau hypersurfaces $\mathcal{P}(z_1, \dots, z_{n+2})$ is fixed under the full $\mathbf{Z}_{n+2}^n$ invariance. We will therefore consider points of $\mathcal{Q}$ that are fixed under subgroups $G_{[\alpha]}$ of $\mathbf{Z}_5^3$. Then we describe the various fractional branes living at these singularities, the corresponding quiver diagrams and the massless chiral fields of the effective theory on the D branes.

As there are several subgroups $G_{[\alpha]}$ in $\mathbf{Z}_5^3$, we shall fix our attention on two categories of subsymmetries; those isomorphic to $\mathbf{Z}_5$; i.e $G_{[1]} \simeq \mathbf{Z}_5$ and those isomorphic to $\mathbf{Z}_5^2$; i.e $G_{[2]} \simeq \mathbf{Z}_5^2$. The Calabi-Yau charges will be taken as in eqs(3.5). Generalization to subgroups of $\mathbf{Z}_{n+2}^n$, though tedious, is a priori straightforward.

7.1 Fixed subspaces of $\mathcal{Q}^{[\alpha]}$

We will consider first the spaces $\mathcal{S}_{(a)}$ of fixed points under a generic $\mathbf{Z}_5$ factor of $\mathbf{Z}_5^3$. Then we examine the spaces $\mathcal{S}_{(ab)}$ of fixed points under $\mathbf{Z}_5^2$ factors. To have an idea on what

these spaces look like, it is interesting to think about the quintic homogeneous hypersurface eq(3.1) as a fiber bundle described by the following equation;

$$P(z_1, \dots, z_5) = \sum_{n_{m+1} \dots n_{m+5}=0}^5 b_{n_{m+1} \dots n_{m+5}} z_{i_{m+1}}^{n_{m+1}} \dots z_{i_{m+5}}^{n_{m+5}}, \quad (7.1)$$

where $b_{n_{m+1} \dots n_{m+5}} = b_{n_{m+1} \dots n_{m+5}}(z_{i_1}, \dots, z_{i_m})$ are some given monomials, in the $z_{i_1}, \dots, z_{i_m}$ complex variables, with appropriate degrees. Let us give examples on how this works in practice.

7.1.1 $\mathbf{CP}^2 \bowtie \mathcal{S}_1$ Fibration

A simple example of realizing fibrations of the quintic consists to rewrite eq(7.1) as,

$$P(z_1, \dots, z_5) = b_{00} + b_{11} z_1 z_2 + b_{50} z_1^5 + b_{05} z_2^5, \quad (7.2)$$

where the b_{mn} coefficient functions are given by

$$\begin{aligned} b_{00}(z_3, z_4, z_5) &= z_3^5 + z_4^5 + z_5^5, \\ b_{11}(z_3, z_4, z_5) &= a_0 z_3 \cdot z_4 \cdot z_5, \\ b_{50}(z_3, z_4, z_5) &= 1, \\ b_{05}(z_3, z_4, z_5) &= 1, \end{aligned} \quad (7.3)$$

and all remaining others are equal to zero. Eqs(7.2-3) mean that the quintic may be viewed as a fibration space whose base $\mathbf{B}_1$ is just the $\mathbf{CP}^2$ space and whose complex one dimension fiber $\mathbf{F}$, to which we refer here below to as $\mathcal{S}_1$, is given by,

$$z_1^5 + z_2^5 + b_1 z_1 z_2 = 0. \quad (7.4)$$

This relation is invariant under the change $(z_1, z_2) \rightarrow (\omega z_1, \omega^{-1} z_2)$; that is under the $\mathbf{Z}_5$ subsymmetry of charges q_i^1 ; the $\mathbf{B}_1$ base space is not viewed at all under this change. The symmetry of the $\mathcal{S}_1$ fiber has one fixed point namely (0,0) and so $\mathcal{S}_1$ is singular at the origin $z_1 = z_2 = 0$. To see what eq(7.4) represents, note that from the z_1 and z_2 variables, one can build three invariant namely $u = z_1^5$, $v = z_2^5$ and $w = z_1 z_2$ having an $\mathbf{A}_4$ singularity. In terms of the new variables, the equation of the $\mathcal{S}_1$ complex curve reads as

$$\begin{aligned} u + v + b w &= 0 \\ uv &= w^5. \end{aligned} \quad (7.5)$$

Therefore near the fixed point $z_1 = z_2 = 0$, the $\mathbf{Z}_5$ orbifold of the commutative quintic $\mathcal{Q}$ can be then viewed as given by the fiber bundle $\mathbf{CP}^2 \bowtie \mathcal{S}_1$ with a vanishing two cycle at $z_1 = z_2 = 0$. Before going ahead, let us comment briefly the complex resolution of this kind of singularity and give its toric geometry diagram representation as shown here below.

$\mathbf{A}_2$ Singularity	$xy = z^2$	(7.6)
$\mathbf{A}_4$ Singularity	$uv = w^5$	
Complex Resolution of $\mathbf{A}_4$	$uv = w^5 + \alpha_4 w^4 + \alpha_3 w^3 + \alpha_2 w^2 + \alpha_1 w + \alpha_0$	
Rules	<i>i)</i> White nodes such as $\bigcirc$, are associated to each non compact $\mathbf{C}$ variables x and y <i>ii)</i> Nodes such as $\otimes$ are associated with blown up spheres with self intersection (-2) <i>iii)</i> Each link $\longleftrightarrow$ represents intersecting spheres with a weight (1)	
Quiver Diagrams	<i>i)</i> Quiver diagram for the resolution of $\mathbf{A}_2$: $\bigcirc_{z^2} \longrightarrow \otimes_z \longleftarrow \bigcirc_1$ <i>ii)</i> Quiver diagram for the resolution of $\mathbf{A}_4$ $\bigcirc_{w^5} \longleftrightarrow \otimes_{w^4} \longleftrightarrow \otimes_{w^3} \longleftrightarrow \otimes_{w^2} \longleftrightarrow \otimes_w \longleftarrow \bigcirc_1$	

More details on this diagram representation of the Kahler and complex resolution of ADE singularities as well as applications in string compactifications may be found in [42-45]

7.1.2 $\mathbf{B} \bowtie \mathcal{S}_{12}$ Fibration

The other example we want to give corresponds to the fibration $\mathbf{B} \bowtie \mathbf{F} \equiv \mathbf{CP}^1 \bowtie \mathcal{S}_{12}$. In this case, the analogue of the above equations read for $\mathcal{S}_{12}$ as follows:

$$P(z_1, \dots, z_5) = b_{000} + b_{111} z_1 z_2 z_3 + b_{500} z_1^5 + b_{050} z_2^5 + b_{005} z_3^5, \quad (7.7)$$

where the b_{mnr} coefficients are follows,

$$\begin{aligned}
b_{000}(z_4, z_5) &= z_4^5 + z_5^5, \\
b_{111}(z_4, z_5) &= a_0 z_4 \cdot z_5, \\
b_{500}(z_4, z_5) &= 1, \\
b_{050}(z_4, z_5) &= 1, \\
b_{005}(z_4, z_5) &= 1,
\end{aligned} \quad (7.8)$$

and all others are equal to zero. The equation of the complex two dimensional singular fiber $\mathcal{S}_{12}$ is;

$$z_1^5 + z_2^5 + z_3^5 + b z_1 z_2 z_3 = 0. \quad (7.9)$$

This is a complex two dimension surface invariant under the change

$$(z_1, z_2, z_3) \rightarrow (\omega^2 z_1, \omega^{-1} z_2, \omega^{-1} z_3); \quad (7.10)$$

that is under the $\mathbf{Z}_5 \otimes \mathbf{Z}_5$ subsymmetry of charges (q_i^1, q_i^2) . The latter does not affect the $\mathbf{B}_{12}$ base space but has one fixed fiber point at $(0, 0, 0)$; so $\mathcal{S}_{12}$ is singular at the origin $z_1 = z_2 = z_3 = 0$. Now introducing the following four invariant $u_1 = z_1^5$, $u_2 = z_2^5$, $u_3 = z_3^5$ and $t = z_1 z_2 z_3$, one sees that they are related as

$$u_1 u_2 u_3 = t^5; \quad (7.11)$$

while the $\mathcal{S}_{12}$ complex surface reads in terms of these invariants as

$$u_1 + u_2 + u_3 + bt = 0.$$

From this relation, one recognizes the two individual singularities associated with each factor of the $\mathbf{Z}_5 \otimes \mathbf{Z}_5$ symmetry. These are given by the following $\mathbf{A}_4$ eqs;

$$\begin{aligned} u_1 u_2 &= \frac{t^5}{u_3}, & \text{for } u_3 \neq 0, \\ u_1 u_3 &= \frac{t^5}{u_2}, & \text{for } u_2 \neq 0. \end{aligned} \quad (7.12)$$

Eq(7.11) describes the case where both of the above singularities collapse; it has a nice description in terms of quivers diagrams.

Singularity	$u_1 u_2 u_3 = t^5$
Complex Resolution	$u_1 u_2 u_3 = t^5 + \alpha_4 t^4 + \alpha_3 t^3 + \alpha_2 t^2 + \alpha_1 t + \alpha_0$
Rules	Same rules as in previous table.
Symmetries	$\mathbf{Z}_5 : \quad z_1 \rightarrow \omega z_1, z_3 \rightarrow \omega^{-1} z_3, \quad z_1 z_3 \text{ is an invariant.}$ $\mathbf{Z}_5 \otimes \mathbf{Z}_5 : z_1 \rightarrow \omega^2 z_1, z_2 \rightarrow \omega^{-1} z_2, z_3 \rightarrow \omega^{-1} z_3,$ $\mathbf{Z}_5 : \quad z_1 \rightarrow \omega z_1, z_2 \rightarrow \omega^{-1} z_2, \quad z_1 z_2 \text{ is an invariant.}$
Quiver Diagram	Here it is represented the three graphs associated to the resolution of the singularities of the discrete symmetries reported above.

(7.13)

Note that the $\mathbf{Z}_5 \otimes \mathbf{Z}_5$ symmetry has a total charge charge $\mathbf{q}^1 + \mathbf{q}^2 = (2, -1, -1, 0, 0)$, behaving then as the $\mathbf{Z}_5$ diagonal symmetry; the remaining off diagonal factor has a total charge $\mathbf{q}^1 - \mathbf{q}^2 = (0, -1, 1, 0, 0)$.

7.1.3 Other Fibrations

Following the same method we have used for $\mathcal{S}_1$ and $\mathcal{S}_{12}$, one can work out the other $\mathbf{B}_a \bowtie \mathcal{S}_a$ and $\mathbf{B}_{(ab)} \bowtie \mathcal{S}_{(ab)}$ Quintic fibrations associated with the natural subgroups of $\mathbf{Z}_5^3$. Denoting the various invariants under the subgroups of $\mathbf{Z}_5^3$ as $u_i = z_i^5$, $w_{ij} = z_i z_j$ and $t_{ijk} = z_i z_j z_k$, one can work out the different equations of the $\mathcal{S}_a$ and fibers $\mathcal{S}_{(ab)}$; the basis $\mathbf{B}_a$ and $\mathbf{B}_{(ab)}$ are respectively given by the $\mathbf{CP}^2$ and $\mathbf{CP}^1$ complex projective spaces. The results are collected in the table eq(7.14).

Fibers	Equations	Fibers	Equations
$\mathcal{S}_1$	$u_1 + u_2 + b_1 w_{12} = 0$ $u_1 u_2 = w_{12}^5.$	$\mathcal{S}_{12}$	$u_1 + u_2 + u_3 + b t_{123} = 0,$ $u_1 u_2 u_3 = t_{123}^5.$
$\mathcal{S}_2$	$u_1 + u_3 + b_2 w_{13} = 0$ $u_1 u_3 = w_{13}^5.$	$\mathcal{S}_{23}$	$u_1 + u_3 + u_4 + b t_{134} = 0,$ $u_1 u_3 u_4 = t_{134}^5.$
$\mathcal{S}_3$	$u_1 + u_4 + b_3 w_{14} = 0$ $u_1 u_4 = w_{14}^5.$	$\mathcal{S}_{13}$	$u_1 + u_2 + u_4 + b t_{124} = 0,$ $u_1 u_2 u_4 = t_{124}^5.$

(7.14)

Having these results at hand, we turn now to give some details by studying the fixed spaces under the orbifold subgroups $\mathbf{Z}_5$ and $\mathbf{Z}_5^2$. We first consider the orbifolds $\mathcal{Q}^{[1]} \simeq \mathcal{R}'/\mathbf{Z}_5$ and then the $\mathbf{Z}_5^2$ orbifolds $\mathcal{Q}^{[2]} \simeq \mathcal{R}''/\mathbf{Z}_5^2$.

These orbifolds corresponds also to start from eq(3.1) and choose either one of the three q_i^a vector charges non vanishing say $q_i^1 = (1, -1, 0, 0, 0)$ while the two others $q_i^2 = q_i^3 = \mathbf{0}$; or two vector charges non vanishing while the third is zero such as, for instance, $q_i^1 = (1, -1, 0, 0, 0)$, $q_i^2 = (1, 0, -1, 0, 0)$ and $q_i^3 = \mathbf{0}$.

7.2 Fractional Branes on $\mathcal{Q}^{[1]}$

As there are three manifest $\mathbf{Z}_5$ subsymmetry factors in the orbifold group $\mathbf{Z}_5^3$, each one generated by an operator F_a , one can write down three corresponding fibrations for the commutative quintic, to which we shall refer hereafter to as $\mathbf{B}_a \bowtie \mathcal{S}_a$. The $\mathbf{B}_a$ spaces are the bases of the fibration and the $\mathcal{S}_a$'s are the fibers encountered above.

7.2.1 Example : $\mathcal{Q}^{[1]} \simeq \mathbf{B}_1 \bowtie \mathcal{S}_1$

Consider the $\mathbf{Z}_5$ subgroup generated by E_1 with q_i^1 charges taken as;

$$q_i^1 = (1, -1, 0, 0, 0), \quad (7.15)$$

the $\mathbf{B}_1 \bowtie \mathcal{S}_1$ fibration of the quintic is just that given by eqs(7.2-4). Since we are working in the coordinate patch $z_5 = 1$, the $\mathbf{B}_1$ base is a patch of $\mathbf{CP}^2$; that is $\mathbf{B}_1 \sim \mathbf{C}^2$ parameterized by the z_3 and z_4 complex coordinates. The codimension two fiber $\mathcal{S}_1$ is given by the following complex curve with an $\mathbf{A}_4$ singularity,

$$u + v + \frac{v^5}{u} = 0 \quad (7.16)$$

7.2.2 First Results

The $\mathbf{B}_a \bowtie \mathcal{S}_a$ fibrations of the quintic are completely determined by the q_i^a Calabi-Yau charges; the $\mathbf{B}_a$ base manifolds are parameterized by those holomorphic coordinates z_i with $q_i^a = 0$ while the $\mathcal{S}_a$ fibers are parameterized by those complex variables with non zero q_i^a

charges with fixed points at the origin. For the q_i^a 's taken as in eq(3.5), we have the following results collected in the table eq(7.17).

Generators F_a	F_1	F_2	F_3
Fixed Points	$(0, 0, z_3, z_4, z_5)$	$(0, z_2, 0, z_4, z_5)$	$(0, z_2, z_3, 0, z_5)$
$\mathbf{B}_a = \mathbf{CP}^2$	$\mathbf{B}_{12} = \{z_3, z_4, z_5\}$	$\mathbf{B}_{23} = \{z_2, z_4, z_5\}$	$\mathbf{B}_{31} = \{z_2, z_3, z_5\}$
$\mathcal{S}_a$ Fibers	$u_1 + w_{12} + \frac{w_{12}^5}{u_1} = 0$ $u_1 = z_1^5, u_2 = z_2^5,$ $w_{13} = z_1 z_3.$	$u_1 + w_{13} + \frac{w_{13}^5}{u_1} = 0$ $u_1 = z_1^5, u_3 = z_3^5,$ $w_{12} = z_1 z_2.$	$u_1 + w_{14} + \frac{w_{14}^5}{u_1} = 0$ $u_1 = z_1^5, u_4 = z_4^5,$ $w_{14} = z_1 z_4.$

(7.17)

Having the above features in mind, the singular representations of the NC quintic may be obtained by starting from the regular representation eqs(5.16) and taking the appropriate limits. For the $\mathcal{S}_1$ singular space for instance, the fields moduli are obtained from eqs(5.16) by setting $q_i^1 = q_i^2 = 0$ and taking the zero limit of z_1 and z_2 . We have for the fiber $\mathcal{S}_1$,

$$\begin{aligned}
\Phi_1 &= (Q \otimes I_{id} \otimes I_{id}) \lim_{z_1 \rightarrow 0} z_1, \\
\Phi_2 &= (Q^{-1} \otimes I_{id} \otimes I_{id}) \lim_{z_1 \rightarrow 0} z_2, \\
F_1 &= P \otimes Q^{\eta_{12}} \otimes Q^{\eta_{13}},
\end{aligned} \tag{7.18}$$

while for $\mathbf{B}_1$, we have:

$$\begin{aligned}
\Phi_3 &= z_3 (I_{id} \otimes I_{id} \otimes I_{id}), \\
\Phi_4 &= z_4 (I_{id} \otimes I_{id} \otimes I_{id}), \\
\Phi_5 &= z_5 (I_{id} \otimes I_{id} \otimes I_{id}), \\
F_2 &= Q^{\eta_{21}} \otimes P \otimes Q^{\eta_{23}}, \\
F_3 &= Q^{\eta_{31}} \otimes Q^{\eta_{32}} \otimes P.
\end{aligned} \tag{7.19}$$

As one sees, once the limit to the singular point is taken, the non vanishing matrix field moduli Φ_i , $i = 3, 4, 5$ are proportional to the identity $(I_{id} \otimes I_{id} \otimes I_{id}) = \mathbf{I}_{\mathcal{D}(G)}$ of the group representation $\mathcal{D}(G)$; i.e $\Phi_i = z_i \otimes \mathbf{I}_{\mathcal{D}(G)}$. This a very remarkable feature at singularity which have an algebraic and a brane interpretations.

Fractional branes on $\mathcal{R}'/\mathbf{Z}_5$

To fix the ideas, let start from the $D9$ brane of type IIB string wrapped on the quintic Q ; and whose coordinates are denoted as $(x^\mu; y_1, y_2, y_3)$ with $x^\mu = (x^0, x^1, x^2, x^3)$ the longitudinal non compact variables of (representing mainly a $D3$ brane embedded in $D9$) and the y_i 's are the compact transverse coordinates of the wrapped $D9$ branes. In other words, the $D9$ brane may naively be thought of as; $D9 \sim D3 \times Q$. In the coordinate patch $z_5 = z_4 = 1$, the y coordinates may be imagined as related to those of the quintic as,

$$\begin{aligned} y_1 &= z_1, \\ y_2 &= z_2, \\ y_3 &= \frac{-a_0 z_1 z_2 z_3}{(2 + z_1^5 + z_2^5 + z_3^5)}, \end{aligned} \tag{7.20}$$

In the case where the orbifold of the quintic is given by $Q^{[1]}$ and torsion taken into account, the wrapped $D9$ becomes a non commutative brane generated by the algebra eqs(6.1) or equivalently by eqs(5.1). At the fixed point of the $Q^{[1]}$ orbifold, where a real two cycle shrinks to zero, the NC wrapped $D9$ becomes a priori a NC wrapped $D7$ described by;

$$\begin{aligned} \Phi_3 &= z_3 I_{\mathcal{D}(\mathbf{Z}_5)}, & \Phi_4 &= z_4 I_{\mathcal{D}(\mathbf{Z}_5)}, & \Phi_5 &= z_5 I_{\mathcal{D}(\mathbf{Z}_5)}, \\ F_1 &= P, & F_2 &= Q^{\eta_{21}}, & F_3 &= Q^{\eta_{31}}. \end{aligned} \tag{7.21}$$

while the singular modes at the orbifold point are carried by the Q operator and its inverse Q^{-1} as shown on the following eqs.

$$\Phi_1 = Q \lim_{z_1 \rightarrow 0} z_1, \quad \Phi_2 = Q^{-1} \lim_{z_2 \rightarrow 0} z_2. \tag{7.22}$$

However due to the complete reducibility property of $I_{\mathcal{D}(\mathbf{Z}_5)}$ namely $I_{\mathcal{D}(\mathbf{Z}_5)} = \sum_{n=1}^5 \Pi_n$, eqs(7.19) shows that they describe rather a set of five commuting Euclidean wrapped $D7$ branes parameterized as

$$\Phi_{3,n} = z_3 \Pi_n, \quad \Phi_{4,n} = z_4 \Pi_n, \quad \Phi_5 = z_{5,n} \Pi_n. \tag{7.23}$$

Therefore at the orbifold point of $\mathcal{R}'/\mathbf{Z}_5$, we have five fractional wrapped $D7$ branes. Moreover since near the singularity the Φ_1 and Φ_2 singular operator may also be split as

$$\Phi_1 = \left(\lim_{z_1 \rightarrow 0} z_1 \right) \sum_{n=1}^5 a_n^+, \quad \Phi_2 = \left(\lim_{z_2 \rightarrow 0} z_2 \right) \sum_{n=1}^5 a_n^-,$$

there are also massless modes $\phi_{1,n} \sim a_n^+$ and $\phi_{2,n} \sim a_n^-$ living on the wrapped $D7$ branes; they are propagating modes traveling between $\Phi_{i,n}$ and $\Phi_{i,n\pm 1}$ fractional branes. The quiver diagram representing the fractional branes is also a Δ_5 pentagon with wrapped $D7$ sitting at the vertices and massless modes represented by the links.

A similar analysis to that we developed above may be also written down for the $\mathbf{B}_2 \bowtie \mathcal{S}_2$ and $\mathbf{B}_3 \bowtie \mathcal{S}_3$ quintic fibrations. The five links joining the neighboring π_n nodes describe massless fields of the effective field theory on the $D4$ branes at singularity. For each fibration, there are (2×5) massless complex fields which we denote as $\chi_{a,k_a} \sim a_{k_a}^+$ and $\psi_{a,k_a} \sim a_{k_a}^-$. In particular at $z_1 = z_2 = z_3 = 0$, we have therefore triplets of massless fields as shown on table eq(7.24) where we give the fields spectrum on the fractional wrapped $D7$ branes on the $\mathbf{B}_a$ basis at the various singularities.

Fields Spectrum

Chiral fields $\rightarrow$	Massless complex fields	Number of massless fields
Singularity $\mathcal{S}_1$	$\chi_{1,k_1}, \psi_{1,k_1}$	2×5
Singularity $\mathcal{S}_2$	$\chi_{2,k_2}, \psi_{2,k_2}$	2×5
Singularity $\mathcal{S}_3$	$\chi_{3,k_3}, \psi_{3,k_3}$	2×5
Locus $z_1 = 0$	$\chi_{1,k_1}, \chi_{2,k_2}, \chi_{3,k_3}$	3×5

(7.24)

Such analysis can also be extended to the case where the orbifold of the quintic is realized as $Q^{[2]} \sim \mathcal{R}''/\mathbf{Z}_5^2$.

7.3 Fractional Branes on $Q^{[2]}$

There are three manifest fibrations of the commutative quintic orbifold $Q^{[2]}$ depending on the nature of the $\mathbf{Z}_5^2$ subgroups of $\mathbf{Z}_5^3$ one is considering. Let us describe them briefly here below

7.3.1 Fibration $Q^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{12}$

Here the $\mathbf{Z}_5^2$ orbifold subsymmetry of the $\mathbf{Z}_5^3$ group has the following charges

$$\begin{aligned}
q_i^1 &= (1, -1, 0, 0, 0), \\
q_i^2 &= (1, 0, -1, 0, 0).
\end{aligned}
\tag{7.25}$$

The q_i^3 charges of the third factor may be thought as being set equal to zero. As such the $\mathbf{B}_{12}$ basis of the fibration is parameterized by the z_4 and z_5 coordinates; while the codimension one fiber $\mathcal{S}_{12}$ is given by the following complex surface with an $\mathbf{A}_4$ singularity,

$$u + v + t + \frac{t^5}{uv} = 0, \quad (7.26)$$

Like for the $\mathbf{B}_a \bowtie \mathcal{S}_a$ fibrations, the $\mathbf{B}_{ab}$ base manifolds are parameterized by those holomorphic coordinates z_i with $q_i^a = 0$ while the $\mathcal{S}_{ab}$ fibers are parameterized by the complex variables with non zero q_i^a charges with fixed points at the origin.

7.3.2 Fibration $\mathcal{Q}^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{23}$

In this fibration, the $\mathbf{Z}_5^2$ orbifold subgroup has the following charges

$$\begin{aligned} q_i^2 &= (1, 0, -1, 0, 0), \\ q_i^3 &= (1, 0, 0, -1, 0). \end{aligned} \quad (7.27)$$

The $\mathbf{B}_{23}$ basis and the $\mathcal{S}_{23}$ fiber of the quintic are respectively parameterized by (z_2, z_5) and (z_1, z_3, z_4) complex variables. Locally $\mathbf{B}_{23} \sim \mathbf{C}$ while the $\mathcal{S}_{23}$ singular surface is of type eq(7.24) with an $\mathbf{A}_4$ singularity at the origin.

7.3.3 Fibration $\mathcal{Q}^{[2]} \simeq CP^1 \bowtie \mathcal{S}_{31}$

In this case, the Calabi-Yau charges of the $\mathbf{Z}_5^2$ orbifold subgroup are;

$$\begin{aligned} q_i^1 &= (1, -1, 0, 0, 0), \\ q_i^3 &= (1, 0, 0, -1, 0). \end{aligned} \quad (7.28)$$

The base is parameterized by the invariant (z_3, z_5) coordinates, while the $\mathcal{S}_{31}$ fiber has a singularity at $z_1 = z_2 = z_4 = 0$. Its complex equation $u_1 + u_2 + t + \frac{t^5}{u_1 u_2} = 0$ has an $\mathbf{A}_4$ singularity at $u_1 = u_2 = t = 0$.

The main features of the various $\mathbf{B}_{(ab)} \bowtie \mathcal{S}_{(ab)}$ fibrations are collected in this table

$F_a \otimes F_b$	$F_1 \otimes F_2$	$F_2 \otimes F_3$	$F_3 \otimes F_1$
Fix Points	$(0, 0, 0, z_4, z_5)$	$(0, z_2, 0, 0, z_5)$	$(0, 0, z_3, 0, z_5)$
$\mathbf{B}_{ab}$	$\mathbf{CP}^1 = \{z_4, z_5\}$	$\mathbf{CP}^1 = \{z_2, z_5\}$	$\mathbf{CP}^1 = \{z_3, z_5\}$
Fibers $\mathcal{S}_{ab}$	$u_1 + u_2 + t + \frac{t^5}{u_1 u_2} = 0$ $u_1 = z_1^5, u_2 = z_2^5,$ $t = z_1 z_2 z_3.$	$u_1 + u_3 + v + \frac{v^5}{u_1 u_2} = 0$ $u_1 = z_1^5, u_3 = z_3^5,$ $v = z_1 z_3 z_4$	$u_1 + u_4 + w + \frac{w^5}{u_1 u_4} = 0$ $u_1 = z_1^5, u_4 = z_4^5,$ $w = z_1 z_2 z_4.$
Fields	$\chi_{1,(k_1,k_2)}, \psi_{1,(k_1,k_2)}$	$\chi_{2,(k_2,k_3)}, \psi_{2,(k_2,k_3)}$	$\chi_{3,(k_1,k_3)}, \psi_{3,(k_1,k_3)}$

(7.29)

The singular representations of non commutative $\mathcal{Q}^{nc[2]}$ may be obtained by starting from the regular representation eqs(5.16) and taking the appropriate limits. For the $\mathcal{S}_{12}$ singular space, the fields moduli are obtained from eqs(5.16) by setting $q_i^1 = q_i^2 = 0$ and taking z_1, z_2 and z_3 to zero. For $\mathcal{S}_{12}$ fiber, we get:

$$\begin{aligned}
\Phi_1 &= (Q \otimes I_{id} \otimes I_{id}) \lim_{z_1 \rightarrow 0} z_1, \\
\Phi_2 &= (Q^{-1} \otimes I_{id} \otimes I_{id}) \lim_{z_2 \rightarrow 0} z_2, \\
\Phi_3 &= (I_{id} \otimes Q^{-1} \otimes I_{id}) \lim_{z_3 \rightarrow 0} z_3, \\
F_1 &= P \otimes Q^{\eta_{12}} \otimes Q^{\eta_{13}}, \\
F_2 &= Q^{\eta_{21}} \otimes P \otimes Q^{\eta_{23}},
\end{aligned} \tag{7.30}$$

while for $\mathbf{B}_{12}$, we have:

$$\begin{aligned}
\Phi_4 &= z_4 \otimes \mathbf{I}_{\mathcal{D}(G)}, \\
\Phi_5 &= z_5 \otimes \mathbf{I}_{\mathcal{D}(G)}, \\
F_3 &= Q^{\eta_{31}} \otimes Q^{\eta_{32}} \otimes P.
\end{aligned} \tag{7.31}$$

Once the limit to the singular point is taken, the non vanishing matrix field moduli Φ_i , $i = 4, 5$ are proportional to the identity $\mathbf{I}_{\mathcal{D}(G)}$ of the group representation $\mathcal{D}(G)$; but as before, this property reflects just the existence of fractional D branes at the orbifold point $z_1 = z_2 = z_3 = 0$.

Fractional branes on $\mathcal{Q}^{[2]}$

Let us reconsider the example of the $D9$ brane of type IIB string wrapped on $\mathcal{Q}^{[2]}$; with local coordinates $(x^\mu; y_1, \dots, y_3)$; the x^μ 's being the longitudinal non compact coordinates of the $D3$ part of $D9$, while the y compact coordinates are as in eqs(7.20).

In case where torsions of $\mathcal{Q}^{[2]}$ are taken into account, the wrapped $D9$ on $\mathcal{Q}^{[2]}$ becomes a non commutative brane generated by the algebra eqs(5.1) where the calabi-Yau charges are chosen as in eqs(7.23). At the fixed point of the $\mathcal{Q}^{[2]}$ orbifold, where a real four cycle shrinks to zero, the NC wrapped $D9$ give rise to twenty five fractional wrapped $D5$ branes on $\mathbf{B}_{12} \sim \mathbf{CP}^1$. The transverse coordinates of these fractional branes are given by;

$$\begin{aligned}\Phi_{4;n,m} &= x_4 \Pi_{n,m}, \\ \Phi_{5;n,m} &= x_5 \Pi_{n,m},\end{aligned}\tag{7.32}$$

where $\Pi_{n,m}$ are the projectors on the $\mathcal{D}(\mathbf{Z}_5^2)$ representation states. The singular modes at the orbifold point are carried by the $Q \otimes Q$, $Q^{-1} \otimes I_{id}$ and $I_{id} \otimes Q^{-1}$ operators of $\mathcal{D}(\mathbf{Z}_5^2)$. Moreover since near the singularity the Φ_1 , Φ_2 and Φ_3 operators may also be split as

$$\begin{aligned}\Phi_1 &= \left(\lim_{z_1 \rightarrow 0} x_1 \right) \sum_{n_1, n_2=1}^5 A_{n_1}^+ A_{n_2}^+, \\ \Phi_2 &= \left(\lim_{z_2 \rightarrow 0} x_2 \right) \sum_{n_1, n_2=1}^5 A_{n_1}^- \Pi_{n_2}, \\ \Phi_3 &= \left(\lim_{z_3 \rightarrow 0} x_3 \right) \sum_{n_1, n_2=1}^5 \Pi_{n_1} A_{n_2}^-, \end{aligned}\tag{7.33}$$

where $A_{n_a}^\pm$ were introduced in subsection eqs(4.30), one gets as a consequence 3×25 massless modes $\phi_{1;n_1,n_2} \sim A_{n_1}^+ A_{n_2}^+$, $\phi_{2;n_1,n_2} \sim A_{n_1}^- \Pi_{n_2}$ and $\phi_{3;n_1,n_2} \sim \Pi_{n_1} A_{n_2}^-$ living on the $D5$ branes wrapped on $\mathbf{CP}^1$; they are propagators between the $\Phi_{i;n_1,n_2}$ and $\Phi_{i,n_1 \pm 1, n_2 \pm 1}$ fractional $D5$ branes. The quiver diagram representing the fractional $D5$ branes wrapped on $\mathbf{CP}^1$ is also a $\Delta_5 \times \Delta_5$ polygon with $D5$ branes sitting at the vertices and the $\phi_{a;n_1,n_2}$ massless modes propagating along the links, see figure 4.

An analogous analysis may be also made for the $\mathbf{B}_{23} \bowtie \mathcal{S}_{23}$ and $\mathbf{B}_{31} \bowtie \mathcal{S}_{31}$ fibrations of the quintic; we get similar results.




Figure 4: Here is represented the 25 vertices of the quiver diagram of the fractional D branes at $x_1 = x_2 = x_3 = 0$ singularity of the $\mathbf{Z}^2$ orbifold subsymmetry of the quintic. The links between the different vertices represents the massless modes on the wrapped $D5$ brane on the base $\mathbf{B}_{12} = \mathbf{CP}^1$

8 Discussion and Conclusion

In this paper we have studied non commutative geometry versus discrete torsion and given an explicit computation of the spectrum of fractional D branes at singularities of Calabi-Yau hypersurface orbifolds $\mathcal{O} = \mathcal{M}/G$. First we have drawn the general setting of this kind of NC geometry and given links with string compactifications on orbifolds with discrete torsion.

Taking the quintic as an illustrating example, we have worked out general and explicit solutions for $\mathcal{Q}^{nc}$, the non commutative extension of the quintic using first a constrained method and then a more involved one baptized as the CP algebra and denoted as $\mathcal{Q} \boxtimes \mathcal{A}_{\mathcal{D}(G)}$. The latter is a fibration whose base is just the commutative orbifold $\mathcal{O}$, which is also the centre of $\mathcal{O} \boxtimes \mathcal{A}_{\mathcal{D}(G)}$, and its fiber is given by $\mathcal{A}_{\mathcal{D}(G)}$, the algebra of the orbifold group representation $\mathcal{D}(G)$. For the case of the quintic, $\mathcal{Q} = \mathcal{O}$, where the coset symmetry $G_{[\alpha]}$ is a subgroup of $\mathbf{Z}_5^3$; that is $G_{[1]} = \mathbf{Z}_5$ or $G_{[2]} = \mathbf{Z}_5^2$, and in presence of a torsion matrix η_{ab} , we have shown that NC geometry is governed by the following features;

- (i) The matrix η_{ab} should be a non symmetric invertible matrix belonging to $SL(3, \mathbf{Z})$.
- (ii) In addition to the classical symmetries of $\mathcal{Q}$ namely invariance under S_5 permutations of the complex variables and the $\mathbf{Z}_5^3$ invariance acting as $z_i \longrightarrow z_i \omega^{q_i^a}$, there is an extra hidden symmetry $\mathbf{Z}_5^3$ induced by torsion, and acting as $z_i \longrightarrow z_i \omega^{p_i^a}$ with p_i^a charges given by

$$p_i^a = \eta_{ab} q_i^b. \quad (8.1)$$

Note that for $\eta_{ab} = 0$ (no torsion), there is no p_i^a 's.

(iii) To each abelian factor $\mathbf{Z}_5 = \{\omega_i; \omega_i^5 = I_{id}\}$ of the geometric symmetry $\mathbf{Z}_5^3$ of the quintic, it is associated an automorphism symmetry $\mathbf{Z}_5^{aut}$ rotating the ω_i elements of $\mathbf{Z}_5$. These two groups do not commute; $\mathbf{Z}_5 * \mathbf{Z}_5^{aut} \neq \mathbf{Z}_5^{aut} * \mathbf{Z}_5$ and were shown to describe a Fuzzy torus $\mathcal{T}_\kappa^2$ with κ , a given fifth root of unity, say $\kappa = \omega = \exp i\frac{2\pi}{5}$. This feature shows that the effective coset symmetry of the orbifold of the quintic is $(\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3$ rather than $\mathbf{Z}_5^3$. Moreover the solutions for the $\mathbf{Z}_i$ generators of the NC quintic are shown to depend on the $\mathcal{D}\left((\mathbf{Z}_5 * \mathbf{Z}_5^{aut})^3\right)$ group realization and can be thought of as describing a five dimensional Fuzzy torus $\mathcal{T}_{\beta_{ij}}^5$, whose β_{ij} cocycle parameters are given by

$$\beta_{ij} = \omega^{\sum_{a=1}^3 (p_i^a q_j^a - p_j^a q_i^a)} \quad (8.2)$$

Such results extend to any complex n dimension hypersurface $\mathcal{H}_n$, $z_1^{n+2} + \dots + z_{n+2}^{n+2} + a_0 \prod_{i=1}^{n+2} z_i = 0$, with orbifold group $\mathbf{Z}_{n+2}^n$. In this case the full symmetry is $(\mathbf{Z}_{n+2} * \mathbf{Z}_{n+2}^{aut})^{\otimes n}$, the torsion matrix η_{ab} should be invertible and non symmetric matrix belonging to $SL(n, \mathbf{Z})$. The corresponding non commutative geometry is mainly given by a real $(n+2)$ dimensional Fuzzy torus $\mathcal{T}_{\beta_{ij}}^{n+2}$ with β_{ij} cocycles as

$$\beta_{ij} = \prod_{a=1}^3 \exp \left[-i \frac{2\pi}{n} (\eta_{ab} - \eta_{ba}) q_i^a q_j^b \right], \quad (8.3)$$

where now the n vectors $\mathbf{q}^a = (q_i^a)$ have $(n+2)$ integer entries; they satisfy the Calabi-Yau condition $\sum_{i=1}^{n+2} q_i^a = 0$ and, for simple and explicit applications, may be chosen as,

$$\begin{aligned} \mathbf{Z}_{n+2} &: \quad \mathbf{q}^1 = (1, -1, 0, 0, \dots, 0, 0, 0), & (1) \\ \mathbf{Z}_{n+2} &: \quad \mathbf{q}^2 = (1, 0, -1, 0, \dots, 0, 0, 0), & (2) \\ &\dots & \\ \mathbf{Z}_{n+2} &: \quad \mathbf{q}^{n-1} = (1, 0, 0, 0, \dots, -1, 0, 0), & (n-1) \\ \mathbf{Z}_{n+2} &: \quad \mathbf{q}^n = (1, 0, 0, 0, \dots, 0, -1, 0). & (n) \end{aligned} \quad (8.4)$$

Because of the richer symmetry of $\mathcal{H}_n = \{z_1^{n+2} + \dots + z_{n+2}^{n+2} + a_0 \prod_{i=1}^{n+2} z_i = 0\}$, the study of fractional branes at orbifold points of $\mathcal{H}_n/G_{[\alpha]}$ is also rich and depends on the orbifold subgroups $G_{[\alpha]}$ of $\mathbf{Z}_{n+2}^n$ one is considering. There are several subgroups in $\mathbf{Z}_{n+2}^n$, the natural ones are the manifest factors $G_{[\alpha]} = \mathbf{Z}_{n+2}^\alpha$, with $1 \leq \alpha \leq (n-1)$. Let us give some comments regarding this point, but forget for a while about string applications by letting n to be a generic

positive integer greater than two and suppose that we have a p -brane, $p > n$, wrapping the $\mathcal{H}_n$ compact manifold.

Fractional branes on $\mathcal{H}_n/\mathbf{Z}_{n+2}$

Since there are n manifest $\mathbf{Z}_{n+2}$ subsymmetries of type eqs(8.4-1,8.4-n) in $\mathbf{Z}_{n+2}^n$, we have n classes of fractional $(p-2)$ -branes at the n kinds of singularities $z_1 = z_{a+1} = 0$. Extending the analysis we have made for the quintic to generic $\mathcal{H}_n$'s by thinking about $\mathcal{H}_n/\mathbf{Z}_{n+2}$ as a fiber bundle $\mathbf{B}_a \bowtie \mathcal{S}_a$ of base $\mathbf{B}_a = \mathbf{CP}^{n-1}$ and a fiber $\mathcal{S}_a$ given by the complex curve $u + v + \frac{v^{n+2}}{u} = 0$ with an $\mathbf{A}_{n-1}$ singularity at the origin $u = v = 0$, we will have d_1 fractional branes at each orbifold point, with $d_1 = \dim(\mathcal{D}(\mathbf{Z}_{n+2}))$, and $2n$ massless modes living on the $(p-2)$ branes. Points in this NC geometry are represented by polygons Δ_{n+2} with $(n+2)$ vertices and $(n+2)$ edges.

Fractional branes on $\mathcal{H}_n/\mathbf{Z}_{n+2}^2$

Before giving the result concerning $\mathcal{H}_n/\mathbf{Z}_{n+2}^2$, note that the action of $\mathbf{Z}_{n+2}^2$ on the z_i variables is additive; that is if one performs two successive $\mathbf{Z}_{n+2}$ actions with different Calabi-Yau charges, say q_i^a and q_i^b respectively, then the total $\mathbf{Z}_{n+2}^2$ action has the charge $q_i^a + q_i^b$,

$$\begin{aligned} \mathbf{Z}_{n+2} & : & z_i & \longrightarrow z_i \omega^{q_i^a}, \\ \mathbf{Z}_{n+2} & : & z_i & \longrightarrow z_i \omega^{q_i^b}, \\ \mathbf{Z}_{n+2}^2 & : & z_i & \longrightarrow z_i \omega^{(q_i^a + q_i^b)}. \end{aligned} \tag{8.5}$$

From these relations, one sees that all happen as if one has only the $\mathbf{Z}'_{n+2}$ diagonal subsymmetry of $\mathbf{Z}_{n+2}^2$ with $(q_i^a + q_i^b)$ charge; the off diagonal subsymmetry, with $(q_i^a - q_i^b)$, is not taken into account. For the example of the $\mathbf{Z}_{n+2}^2$ symmetry associated to eq (8.4-1) and eq(8.4-2), the diagonal $\mathbf{Z}'_{n+2}$ symmetry has the following vector charge

$$\mathbf{Z}_{n+2}^2 : \quad \mathbf{q}^1 + \mathbf{q}^2 = (2, -1, -1, 0, \dots, 0, 0, 0), \tag{8.6}$$

and fix points at $z_1 = z_2 = z_3 = 0$ and more generally at $z_1 = z_{a+1} = z_{b+1} = 0$. This property shows that:

(a) the fibration $\mathbf{B}_{ab} \bowtie \mathcal{S}_{ab}$ of the manifold $\mathcal{H}_n$ has as a base $\mathbf{B}_{ab} = \mathbf{CP}^{n-2}$ and as a fiber $\mathcal{S}_{ab}$ the complex surface $u + v + t + \frac{t^{n+2}}{uv} = 0$ with an $\mathbf{A}_{n-1}$ singularity at the origin $u = v = t = 0$.

(b) there are $\frac{n(n-1)}{2}$ different fibrations of $\mathcal{H}_n$ and so there are $\frac{n(n-1)}{2}$ classes of fractional $(p-4)$ -branes at the $z_1 = z_{a+1} = z_{b+1} = 0$ singularities. Each class contains $d_1 d_2$ fractional

$(p-4)$ -branes. Using a similar analysis to that of subsections 7.2-3, one can check that there are $2(d_1 + d_2)$ massless modes living on these branes, with $d_1 + d_2 = \dim \mathcal{D}(\mathbf{Z}_{n+2}^2)$. Points in this NC geometry are given by the crossed product of two polygons $\Delta_{d_1} \times \Delta_{d_2}$.

Fractional branes on $\mathcal{H}_n/\mathbf{Z}_{n+2}^k$

Extending the above reasoning to $\mathcal{H}_n/\mathbf{Z}_{n+2}^k$, the orbifold symmetry $\mathbf{Z}_{n+2}^k$ with charges $\mathbf{q}^{a_1}, \dots, \mathbf{q}^{a_k}$ acts in practice through its diagonal $\mathbf{Z}_{n+2}$ subgroup with a Calabi-Yau charge $Q_{(a_1, \dots, a_k)} = \mathbf{q}^{a_1} + \dots + \mathbf{q}^{a_k}$; it has fix points located at $z_1 = z_{a_1+1} = \dots = z_{a_k+1} = 0$. For the example of the leading k symmetries of eqs(8.4), the corresponding diagonal symmetry charge $Q_{(1, \dots, k)}$ is given by;

$$\mathbf{Z}_{n+2}^k : \quad Q_{(1, \dots, k)} = (k, -1, -1, \dots, -1, 0, \dots, 0, 0); \quad (8.7)$$

where the first $(k+1)$ entries of $Q_{(1, \dots, k)}$ are non zero while the remaining ones are all of them equal to zero. A simple counting shows that there are $\frac{n!}{(n-k)!k!}$ possible subgroups of this kind in $\mathbf{Z}_{n+2}^n$.

The fibration $\mathbf{B}_{(a_1 \dots a_k)} \bowtie \mathcal{S}_{(a_1 \dots a_k)}$ of the manifold $\mathcal{H}_n$ has as a base $\mathbf{B}_{(a_1 \dots a_k)} = \mathbf{CP}^{n-k}$ and as a fiber $\mathcal{S}_{(a_1 \dots a_k)}$ the complex dimension k hypersurface $u_1 + \dots + u_k + t + \frac{t^{n+2}}{\varrho} = 0$, $\varrho = \prod_{j=1}^k u_j$, with an $\mathbf{A}_{n-1}$ singularity at $u_1 = \dots = u_k = t = 0$. An other result is that there are $\frac{n!}{(n-k)!k!}$ different fibrations of $\mathcal{H}_n$ and so there are $\frac{n!}{(n-k)!k!}$ classes of fractional $(p-2k)$ -branes at the $u_1 = \dots = u_k = t = 0$ singularities. Each class contains $\left(\prod_{j=1}^k d_j\right)$ fractional $(p-2k)$ -branes $k \sum_{i=1}^k d_i$ massless modes living on these branes, with $\sum_{i=1}^k d_i$ being the dimension of the $\mathcal{D}(\mathbf{Z}_{n+2}^k)$ representation group of $\mathbf{Z}_{n+2}^k$.

The analysis we made for the quintic extends to all homogeneous hypersurfaces and the results we have obtained are also valid. For the special case $n=4$, we recover known results on orbifolds of the $\mathbf{K}_3$ surface. In future work, we will discuss the case NC local Calabi-Yau manifolds embedded in NC toric varieties.

Acknowledgments

This research work is supported by DFG/CNCPRST 445 Mar 113/5/0 contract between University of Munich, Germany, and University of Rabat, Morocco. I am very grateful to Julius Wess for kind hospitality at Munich University, to DFG program which supported my stay in Germany and to CNR Morocco for covering my Air faire. I thank I. Benkaddour, C. Blooman for discussions and A.Belhaj and M.Bennai for earlier collaboration on this subject. Many thanks to Mrs M.Kurzinger and to the team of the theoretical Physics department of Munich

university who made my stay in Germany comfortable.

9 References

References

- [1] David Berenstein, Vishnu Jejjala, Robert G. Leigh, D-branes on Singularities: New Quivers from Old, Phys.Rev. D64 (2001) 046011, hep-th/0012050.
- [2] David Berenstein, Robert G. Leigh, Phys.Lett. B499 (2001), hep-th/0009209.
- [3] Matthias Klein, Raul Rabadan, $Z_N \times Z_M$ orientifolds with and without discrete torsion, JHEP 0010 (2000) 049, hep-th/0008173.
- [4] Eric R. Sharpe, Recent Developments in Discrete Torsion, Phys.Lett. B498 (2001) 104-110, hep-th/0008191.
- [5] Paul S. Aspinwall, M. Ronen Plesser, D-branes, Discrete Torsion and the McKay Correspondence, JHEP 0102 (2001) 009, hep-th/0009042.
- [6] Paul S. Aspinwall, A Note on the Equivalence of Vafa's and Douglas's Picture of Discrete Torsion, JHEP 0012 (2000) 029, hep-th/0009045.
- [7] Keshav Dasgupta, Seungjoon Hyun, Kyungho Oh, Radu Tatar, Conifolds with Discrete Torsion and Noncommutativity, JHEP 0009 (2000) 043, hep-th/0008091.
- [8] M.R Douglas, D-branes and Discrete torsion, hep-th/9807235.
- [9] M.R Douglas and B.Fiol, D-branes and Discrete torsion, hep-th/9903031.
- [10] David Berenstein, Robert G. Leigh, Discrete torsion, AdS/CFT and Duality, JHEP 01 (2000) 038, hep-th/0001055.
- [11] A. Connes, M.R. Douglas et A. Schwarz, JHEP 9802, 003 (1998), hep-th/9711162.
- [12] N. Seiberg and E. Witten, JHEP 9909(1999) 032, hep-th/990814.
- [13] David J. Gross, Nikita A. Nekrasov, Solitons in Noncommutative Gauge Theory, 0103 (2001) 044 hep-th/0010090.
- [14] I. Benkaddour, M. Bennai, E. Diaf and H. Saidi CQG 17(2000)1765.

- [15] A. Belhaj, M. Hssaini, E. L. Sahraoui, E. H. Saidi, *Class.Quant.Grav.* **18**(2001) 2339-2358, hep-th/0007137.
- [16] N. Nekrasov and A. Schwarz, *Commun Math. Phys* **198**(1998) 689-703, hep-th/9802068.
- [17] David Berenstein, Robert G. Leigh, Resolution of Stringy Singularities by Non-commutative Algebras, *JHEP* 0106 (2001) 030, hep-th/0105229.
- [18] M. Bertolini, P. Di Vecchia, M. Frau, A. Lerda, R. Marotta, N=2 Gauge theories on systems of fractional D3/D7 branes, *Nucl.Phys. B* **621** (2002) 157-178, hep-th/010705.
- [19] M. Cvetič, G.W. Gibbons, James T. Liu, H. Lu, C.N. Pope, A New Fractional D2-brane, G_2 Holonomy and T-duality, hep-th/010616.
- [20] Tadashi Takayanagi, Holomorphic Tachyons and Fractional D-branes, *Nucl.Phys. B* **603** (2001) 259-285, hep-th/0103021.
- [21] Subir Mukhopadhyay, Koushik Ray, Fractional Branes on a Non-compact Orbifold, *JHEP* 0107 (2001) 007, hep-th/0102146.
- [22] M. Frau, A. Liccardo, R. Musto, The Geometry of Fractional Branes, *Nucl.Phys. B* **602** (2001) 39-60, hep-th/0012035.
- [23] M. Billo', L. Gallot, A. Liccardo, Fractional branes on ALE orbifolds , the proceedings of the RTN meeting "The Quantum Structure of Spacetime and the Geometric Nature of Fundamental Interactions", (Corfu, September 2001) , hep-th/0112190 .
- [24] A. Belhaj, E.H. Saidi, On Non Commutative Calabi-Yau Hypersurfaces, hep-th/0108143, To appear in *Phy. Let.B*.
- [25] I. Bars, H. Kajiura, Y. Matsuo, T. Takayanagi, Tachyon Condensation on Noncommutative Torus, *Phys.Rev. D* **63** (2001) 086001, hep-th/0010101.
- [26] B. R. Greene, C. I. Lazaroiu, Piljin Yi, D-particles on T^4/Z_n orbifolds and their resolutions, hep-th/9807040; *Nucl.Phys. B* **539** (1999) 135-165.
- [27] E. M. Sahraoui, E.H. Saidi, Solitons on compact and noncompact spaces in large noncommutativity, hep-th/0012259, *Class.Quant.Grav* (2001).
- [28] E. M. Sahraoui, E.H. Saidi, D-branes on Noncommutative Orbifolds, hep-th/0105188.

- [29] Julius Wess and J Bagger, SUPERSYMMETRY AND SUPERGRAVITY, Princeton Univ. Press, (1983).
- [30] Juan M. Maldacena, The Large N Limit of Superconformal Field Theories and Supergravity, Adv.Theor.Math.Phys. 2 (1998) 231-252; Int.J.Theor.Phys. 38 (1999) 1113-1133 , hep-th/9711200 .
- [31] Edward Witten, Multi-Trace Operators, Boundary Conditions, And AdS/CFT Correspondence, hep-th/0112258.
- [32] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri, Y. Oz, Large N Field Theories, String Theory and Gravity, Phys.Rept. 323 (2000) 183-386 ,hep-th/9905111 .
- [33] Non Commutative Local Calabi-Yau manifolds with torsion, In preparation:
- [34] S. Gukov, C. Vafa, E. Witten, CFT's From Calabi-Yau Four-folds, Nucl.Phys. B584 (2000) 69-108, hep-th/9906070.
- [35] Kentaro Hori, Hiroshi Ooguri, Cumrun Vafa, Non-Abelian Conifold Transitions and N=4 Dualities in Three Dimensions, Nucl.Phys. B504 (1997) 147-174, hep-th/9705220.
- [36] Julius Wess, Proceedings of the Workshop on High Energy Physics 2 (2000)1-11. Rabat University.
- [37] Philip Candelas, Eugene Pervalev, Govindan Rajesh, Toric Geometry and Enhanced Gauge Symmetry of F-Theory/Heterotic Vacua, Nucl.Phys. B507 (1997) 445-474, hep-th/9704097.
- [38] A. Lawrence, N. Nekrasov, C. Vafa, On Conformal Theories in Four Dimensions, Nucl.Phys. B533 (1998) 199-209, hep-th/9803015.
- [39] A. Belhaj, E.H.Saidi, Hyperkahler Singularities in Superstrings Compactification and 2d N=4 Conformal Field Theory, Class.Quant.Grav. 18 (2001) 57-82,hep-th/0002205.
- [40] Branislav Jurco, Lutz Möller, Stefan Schraml, Peter Schupp, Julius Wess, Eur.Phys.J. C21 (2001) 383-388, hep-th/0104153.
- [41] John Madore, Stefan Schraml, Peter Schupp, Julius Wess, Gauge Theory on Noncommutative Spaces, Eur.Phys.J. C16 (2000) 161-167, hep-th/0001203.

- [42] Sheldon Katz, David R. Morrison, M. Ronen Plesser, Enhanced Gauge Symmetry in Type II String Theory, Nucl.Phys. B477 (1996) 105-140, hep-th/9601108.
- [43] S.Katz, P.Mayr, C.Vafa, Mirror symmetry and Exact Solution of 4D N=2 Gauge Theories I, Adv.Theor.Math.Phys. 1 (1998) 53-114, Adv.Theor.Math.Phys. 1 (1998) 53-114.
- [44] N.C.Leung, C.Vafa, Branes and Toric Geometry, Adv.Theor.Math.Phys.2 (1998) 91-118, hep-th/9711013.
- [45] A. Belhaj, A. E. Fallah , E. H, Saidi, CQG 17 (2000)515-532.